


CURVATURES AND DISEASE

OF THE SPINE

BRODHURST



Glass Case

For Ref. only

SC WE BRO

C

H. Z. Baker Esq.

with kind regards

W. H. W.

May 1883

ON
CURVATURES AND DISEASE
OF
THE SPINE



Digitized by the Internet Archive
in 2014

<https://archive.org/details/b21289888>

ON
CURVATURES AND DISEASE
OF
THE SPINE

BY

BERNARD E. BRODHURST, F.R.C.S.

FOREIGN ASSOCIATE OF THE ACADEMY OF SCIENCES OF ROME;
CORRESPONDING MEMBER OF THE IMPERIAL SOCIETY OF SURGERY OF PARIS, AND OF THE MEDICAL
AND SURGICAL SOCIETIES OF ROME, LYONS, AND ODESSA;
SURGEON TO THE ROYAL ORTHOPÆDIC HOSPITAL, AND TO THE ROYAL HOSPITAL FOR INCURABLES;
CONSULTING SURGEON TO THE BELGRAVE HOSPITAL FOR CHILDREN;
LATE SURGEON TO THE ORTHOPÆDIC DEPARTMENT, AND LECTURER ON ORTHOPÆDIC SURGERY
AT ST. GEORGE'S HOSPITAL.

THIRD EDITION



LONDON
J. & A. CHURCHILL
NEW BURLINGTON STREET
1883

1871

958026

PREFACE

TO THE

THIRD EDITION

The following Chapters were written as Lectures, which were delivered in the School at St. George's Hospital. In preparing this Edition some additions have been made which I believe will increase any value which may attach to the volume. I have endeavoured to reduce rather than to expand the volume; and to place the subject matter plainly and concisely before the reader. I consider it both undesirable and useless to introduce exploded theories, which have nothing to recommend them but their antiquity, simply for the sake of disproving them. And thus I have contented myself with presenting my subject in as few words as I could employ to render it intelligible.

B. E. B.

January, 1883.

PREFACE

TO THE

SECOND EDITION

THE subject of Spinal Curvatures has not perhaps received all that attention which it deserves. Sir Benjamin Brodie, in a lecture which he delivered at St George's Hospital, in 1846, for instance, says, "When I first became engaged in a considerable private practice, and cases of Curvature of the Spine were presented to my observation, I was in doubt as to their nature and treatment. I knew nothing of them from my own experience, and I had learned nothing of them from my teachers. I felt that in this respect my education had been imperfect, and it took me no small trouble to supply the deficiency. Your situation," continued Sir Benjamin, "in this respect is probably not very different from what mine was formerly." This is a sad reproach, and yet I would ask, has anything been done to remove it. Has the student of 1864 opportunities of studying these affections which were not enjoyed at the time Sir Benjamin Brodie was speaking? I fear it must be said at the present day, as it was said twenty years ago, "that the hospital (and hospitals in general) will not afford the information which we stand in need of." Should this work be useful in teaching

something of the nature and treatment of spinal curvatures, I shall be well pleased to have spent time in writing it.

This Second Edition has been entirely re-written ; it has been much enlarged, and all the woodcuts are new. I am indebted to Mr. Bagg for the great care and accuracy with which he has made the drawings. They were made with the *camera lucida*, and show very truly and exactly what they are intended to illustrate. I am also indebted to my friend Mr. Wheeler, of Bexley, for permission to use several very important pathological specimens in illustration of the subject.

Throughout it has been my endeavour to adhere as strictly to truth as possible. I have desired not to exaggerate the importance of my subject, and yet to state it so fully as to call attention to a class of diseases which is at present neglected. No one can be better aware that I am of the imperfections of this volume ; but if my professional brethren will give me credit for a sense of duty in its preparation, and will accord to this Edition not less favour than to the first, I shall be well rewarded.

20, GROSVENOR STREET ;
October, 1864.

CONTENTS

CURVATURES OF THE SPINE

	PAGE
INTRODUCTORY	1
CHAPTER I	
LATERAL CURVATURE OF THE SPINE	9
CHAPTER II	
FORMATION OF SPINAL CURVES	24
CHAPTER III	
TREATMENT	49
CHAPTER IV	
ANTERIOR CURVATURE OF THE SPINE	66
CHAPTER V	
POSTERIOR CURVATURE OF THE SPINE	73

DISEASE OF THE SPINE

CHAPTER VI	
DISEASE OF THE SPINE AND ANGULAR CURVATURE	78
CHAPTER VII	
DIAGNOSIS AND PROGNOSIS	102
CHAPTER VIII	
TREATMENT	108

CURVATURES OF THE SPINE

INTRODUCTORY

CURVATURES of the spine are known as anterior curvature (*lordosis*), posterior curvature (*cyphosis*), and lateral curvature (*scoliosis*). These terms—lordosis, cyphosis, and scoliosis—have been applied to curvatures of the spine ever since the time of Hippocrates. They were not used then, however, exactly in the same sense as they are now understood; for the term scoliosis, which now signifies lateral curvature of the spine, was then frequently applied indiscriminately to every form of spinal distortion.

Antero-posterior pathological curves are, for the most part, merely exaggerated conditions of the physiological curves. Thus we find anterior curvature existing especially in the lumbar region, and posterior curvature in the dorsal region. It will be shown farther on, however, that these conditions are not absolutely limited to a given region; but that the vertebræ above and below may be involved in the curve, so as to reverse the normal curve of any region. These curves are rarely congenital; but they are for the most part induced through ignorance or neglect on the part of the nurse or through deficient nutrition.

Before proceeding to consider these pathological conditions, however, it may be desirable to examine the normal—physiological—curves of the spinal column,

and to consider very briefly their formation and their purposes.

The spine in the foetus is bent forward; its curve being moulded by the walls of the uterus: the head is bent upon the breast, and the thighs are folded upon the abdomen.

In the infant, the spine remains almost straight while in the horizontal position; the sacrum is only slightly curved, and the pelvis is more oblique than in the adult.

The normal—antero-posterior—curves are developed slowly, and depend in great measure on muscular action and on the erect position which man is destined to assume. At birth these curves do not exist; and even in the young child they are not constant, for they disappear in the horizontal position. Gradually, however, the antero-posterior curves become more permanent; and, long before growth is complete, the spine is found to have assumed, as a constant condition, lumbar, dorsal, and cervical curves; while, on the other hand, it has lost much of its flexibility.

On examining the spinal column of the adult in the erect position, it will be seen that the base of the sacrum is so placed that it is to be found immediately above a straight line which may be supposed to pass through the heads of the thigh-bones. This is known as the inter-femoral line. It is seen in Fig. 1.

Naegele first showed exactly the position of the pelvis in the upright posture. Prior to his demonstration of the fact, it was believed that the pelvis was more horizontal than is absolutely the case. The pelvis is so oblique that the anterior wall (pubes) presents upwards and backwards, and the posterior wall (sacrum and coccyx) downwards and forwards, as is shown in Fig. 2. Such, then, being the positions of the pelvis and of the sacro-lumbar articulation, the question arises, how is equilibrium maintained?

Equilibrium is the result of certain forces acting in front of and behind the spinal column; namely, the action

FIG. 1.





FIG. 2.



of the flexors and the extensors of the trunk, neck and head, and the resistance of the ligaments that hold together the several vertebræ which form the spinal column. In the erect posture these structures combine to poise the head and the vertebral column in the vertical line on the heads of the thigh-bones. (See Fig. 3).


In considering this question, the weight of the viscera should not be forgotten. At the present time, however, the chief points alone can be taken into consideration.

The position of the sacro-vertebral articulation being, then, such as has been described, it becomes necessary for the maintenance of equilibrium, and that the head may be poised (its centre of gravity corresponding with a vertical line which coincides with the axis of the trunk, and which shall fall on the interfemoral line), *first, that the lumbar vertebræ shall be curved backwards.* This lumbar curve is the result of muscular action in the endeavour to maintain the erect position. It is the reverse of the sacral curve; and it springs from the sacro-lumbar articulation upwards and backwards, the sacral curve commencing at the same articulation, and presenting downwards and forwards. When a curve has been formed, it is essential that a second curve—dorsal—shall restore the direction of the spine towards the perpendicular line; and, for a similar reason, a cervical curve is formed in the same direction as the lumbar curve. These several portions of the spine differ somewhat in flexibility, the lumbar and the dorsal portions possessing this quality very nearly in equal degrees; but in the cervical region the flexibility is three times greater than in either of the other regions. Thus, a structure is composed of segments of various circles, strong, yet elastic, fitted to bear the superincumbent weight, and yet capable of resisting the effects of shock.

It would, perhaps, be impossible to imagine a structure more perfectly adapted to its purposes than the spinal column, formed as it is of separate pieces, which are severally united by discs of fibro-cartilage, bound together by numerous and strong ligaments, and surrounded and

acted on by powerful muscles. Its flexibility and its curved form are, perhaps, of more importance even than

FIG. 3.




its great strength: these adapt the spine specially to its purposes.

I do not desire in this place to enter minutely into questions of physiology, and will therefore only allude to the opinions of some eminent physiologists, such as the Webers, Cruveilhier, Bishop, and others, who have stated their belief that the normal curves of the spine depend on varying thicknesses of the bodies of the vertebræ and of the intervertebral substances. It is true, as the Webers have shown, that, in the dorsal region of the spine especially, differences in thickness exist on the anterior and the posterior surfaces of the vertebræ; namely, that the vertebræ are somewhat thinner on their anterior surfaces, in the concavity of the curve, than they are posteriorly, while in the lumbar and in the cervical regions the fibro-cartilages are especially affected being compressed posteriorly.

To show the varying thicknesses of the anterior and the posterior surfaces of the vertebræ and of the intervertebral cartilages, the Webers caused the spine to be bisected in the mesial plane, after the viscera had been removed from the cavities of the thorax and abdomen, the space having been filled up with liquid plaster of Paris. The relative normal positions of the various portions of the spinal column were thus obtained, and from the section the accompanying figure (Fig. 4) was taken, which represents very accurately the forms of the several curves and the proportions of the bones and cartilages which together constitute the spinal column.

But it has been already shown that the physiological—antero-posterior—curves do not exist at birth; and that, when first observed, they are not permanent, but are lost in the horizontal posture. Further, these inequalities in the vertebræ and in the intervertebral substances do not exist at birth, nor are they found until the spinal curves commence to be permanent. Also, they increase with age. Such being the case, it follows as a necessary consequence that these differences in the anterior and the posterior surfaces of the bodies of the vertebræ, and especially in the fibro-cartilages, are due to the antero-posterior curves

FIG. 4.



—are due, in fact, to the erect position and to the pressure from above downwards consequent on that position.

The effect of pressure on the intervertebral cartilages is shown by the loss of height which is sustained at the end of the day. Thus it is well known that a man of middle stature, who remains in the erect posture during the day, will lose nearly one inch in height, and that this is regained only after he has been in the recumbent position for six or eight hours. We must conclude, therefore, that these varying thicknesses of the bodies of the vertebræ and of the intervertebral substances are produced by pressure as a result of the erect position, and that they would not occur, any more than the antero-posterior curves themselves would be developed, in the horizontal position.

Thus, it may be stated that the antero-posterior curves of the spinal column result as a consequence of the erect position and from muscular action in maintaining equilibrium, and that unequal pressure occasions thinning of the bodies of the vertebræ and of the intervertebral substances in the concavities of the curves. This view will be confirmed by the consideration of pathological curves.

CHAPTER I

LATERAL CURVATURE OF THE SPINE

IN lateral curvature of the spine the vertebræ deviate from the mesial line of the trunk. This form of curvature may be divided into two stages, or periods; namely, *incipient* and *confirmed*.

By incipient curvature is understood such a condition of lateral deviation of the spine as is removable in the recumbent position, with the help perhaps of some slight pressure; while a confirmed curve is curvature with rotation of the vertebræ.

Lateral curvature is the most common form of spinal curvature. It occurs more frequently in the female than in the male; it seldom commences after the age of puberty, and it does not commonly occur as a primary affection.

Causes.—The principal predisposing causes are debility—muscular or constitutional, acquired or inherited—and rickets; and the exciting or proximate causes are bad habits of sitting and standing, and through fatigue induced perhaps by overwalking.

The effect of climate is often seen in the debility which precedes curvature of the spine. English children, especially girls, who are born in India, Australia (especially Melbourne), and the West Indies, if they are not sent home early, are extraordinarily liable to flat-foot, knock-knee, and spinal curvature. They are much more liable to these affections than those who are born in northern climates. And it may be stated generally that it is rare for any English girl who is sent home later than eight

years of age to escape flat-foot and spinal curvature. Children become pale and weedy; and even more so in Melbourne than in India. It is probable that many who are brought over suffer through the amount of exercise which they undergo whilst associating with children born in this country. Some at ten years of age are brought over with spinal curvature already developed, and many are brought over from Australia even younger, in whom spinal curvature is advanced. The looser fibre of the southern children will not bear the exercise which is imposed on our more hardy northern children; and it is soon discovered after they arrive that there is something essentially wrong in their manner of standing and walking.

Over-exercise tells immediately on the southern child; it tells prejudicially also on those of this country. There is no country in the world where constitutional walks are imposed on children of tender years as with us: old and young walk alike. At school, those of eighteen and those of ten years of age walk for one hour or for two hours daily as is the rule of the school, quite irrespective of strength. In many of these schools there is not a girl who is not flat-footed. These daily walks are exceedingly injurious to all who are not in robust health. And, indeed, when it becomes a daily parade along the same bit of hard road, without any object to engage the attention or to inspire the desire to cover a certain distance, the result can only be weariness and lassitude. This may be largely noticed at Brighton, for instance, where the melancholy spectacle is to be observed twice daily of girls of all ages made to parade backwards and forwards, two and two, on a concrete walk, without any object but to show themselves, for the benefit of their house. The result of these walks is highly disastrous.

Sitting for prolonged periods in a constrained attitude, such as is often assumed in drawing or in writing; or sitting for a long time on a narrow seat without support for the back will cause the trunk to be inclined to one side or the other. In the first case position favours this

inclination of the trunk, and in the second the trunk is allowed to incline to one side or the other, to obtain relief from the constrained attitude.

Again, the habit of standing, more or less on one leg, induces curvature. This habit is indulged in to a great extent, either through some local weakness or from fatigue; for weariness is readily induced during the period of growth, and especially when growth is rapid. This attitude causes the hip to become prominent, and the pelvis to be inclined towards the opposite side. Together with this obliquity of the pelvis, the spine becomes curved. At first, the curve is temporary; but at length, as the obliquity of the pelvis becomes permanent, the curve also becomes confirmed. Equilibrium is restored by means of secondary, or compensating curves; and these are formed in regular sequence, and are determined by the position of the primary curve.

This is shown in Fig. 5. In this instance through standing on the right foot the pelvis became oblique, falling towards the opposite side, and a lumbar curve was necessarily produced, having its convexity towards the lowest ilium; namely, to the left side of the body.

The following is an outline of the commonest commencement of lateral curvature. A young person, of from 12 to 15 years of age, presents herself with flat-foot, the left foot being flatter than the right, and the left knee perhaps somewhat inverted. The pelvis is inclined towards the left side, so that the left side of the pelvis is sensibly lower than the right. There is a lumbar curve, with its convexity to the left side, and a compensating dorsal curve with its convexity to the opposite side. The right scapula is raised and prominent, and the left shoulder drops.

Figure 6 shows this case reversed. In this instance there was flat-foot and knock-knee more marked on the right than on the left side, so that the right side of the pelvis was lower than the left side. This gave rise to a right lumbar curve, and a compensating left dorsal curve. The

right shoulder was three fourths of an inch lower than the left, and the right side of the pelvis was half an inch lower than the left side.

It is because the left lower extremity is more frequently and more markedly affected than that of the right side, that there is found for the most part a left lumbar curve and a

FIG. 5.



right dorsal curve. A right dorsal curve occurs about nine times more frequently than a left dorsal. And in ninety-five cases in every hundred the lumbar curve is the primary and the dorsal is the secondary curve; that is to say, in these ninety-five cases the cause is from below (flat-foot or knock-knee or both or muscular weakness), occasioning

an oblique pelvis and a lumbar curve, with a consecutive dorsal curve.

I find that of the last 1000 cases of primary lumbar curvature which have come under care in my practice, 893 have been curved to the left, and 107 to the right side.

A delicate child, if treated in the same manner as a

FIG. 6.



stronger companion, will probably develop some irregularity of growth. The particular form of distortion will, however, depend mainly on the habits of the child. Thus, if the child stand or walk much, the ligaments of the ankle-joint and those in the sole of the foot will yield and occasion flat-foot; and the internal lateral ligament of the

knee-joint may yield and cause genu valgum. And if the foot or the knee of one side becomes either flatter or more inverted than on the other side, so as to produce flat-foot or knock-knee, then, by so much as the knee is more inverted and the foot is flatter, the pelvis becomes oblique. But an oblique pelvis must give rise to spinal curvature.

Inequality in length of the lower limbs, whether produced by a bent tibia or femur, by flat-foot or knock-knee, by muscular contraction, articular disease, or by partial loss of muscular power, will induce obliquity of the pelvis and a primary lumbar curve.

It has been stated that the lower limbs are frequently not symmetrical, and that on measuring the two limbs one will be found longer than the other, the difference varying from one eighth of an inch to one inch and a half. Dr Morton, of Philadelphia, measured the limbs of 512 boys and found that in 271 there was inequality of length, and of these the right limb was the longest in 198.

I have failed to verify this statement, and I believe it will be found that in our population this difference does not exist.

If a healthy boy is measured—one who is well grown, and who has never had any bone or joint affection—from the great trochanter to the inner malleolus it is rare to find the least difference in the length of the limbs. It occurs occasionally, but it is generally found that the shorter limb has been badly bruised, or that there has been at least some synovitis, or that the boy has been engaged in exercises beyond his strength.

Even when one of the lower limbs has suffered from infantile paralysis there is frequently no inequality in length. Of course there is great inequality in girth, but frequently not in length. In a paralysed limb there may be shortening to the extent of two inches or more; but this is rare. It is frequently said that there is a short limb when there is only a flat foot; the leg bones and the thigh bones being equal in length on both sides. Selecting healthy boys who have neither been

bruised nor had inflamed joints, probably not one in 500 would show asymmetry.

My attention was called to this subject very specially some years since by one who thought that he was able to account for obliquity of the pelvis through inequality in length of the lower limbs. In examining his cases it was found that in every one of them either one foot was flatter than the other, or that there had been some bone or joint affection. To increase the sole of the boot in thickness for flat-foot alone, is to do an undoubted injury.

Every curvature is produced by a special cause ; and it need scarcely be said that while the cause remains, the curve itself cannot be removed, or if removed it will certainly recur so long as the cause exists. It is of vital importance, therefore, in the treatment of spinal curvature, to ascertain precisely what is the exciting cause in each case of curvature. Until this is done, and the cause is removed, it is useless to attempt to treat a spinal curve, except in the recumbent position.

Again, other causes of spinal curvature exist, namely, such as increase or diminish the power of a single limb. Thus the power may be increased by extraordinary use, as through carrying a burden on one arm, or as occurs in certain trades, where one arm is much more employed than the other. Fig. 7, was taken from a young person who habitually carried a heavy child on the left arm. And it is found that nurses, needlewomen, tailors, shoemakers, compositors, and some others are unusually liable to spinal curvature through the inordinate use of one arm. Also, when muscular power is diminished as through paralysis, partial or complete, of an upper extremity, or after amputation at the shoulder-joint or immediately below the joint, lateral curvature of the spine will occur in the dorsal region, the convexity of the curve forming towards the opposite side. This is shown in Fig. 8, where amputation of the left arm caused the spinal column to be drawn away from the median line and to be curved towards the right side. This drawing was

taken from a powerful man, who after the loss of the left arm continued to use a heavy hammer in the right hand. But, also, it may be that there is little or no increase of muscular power on the side of the convexity, but that the force of pressure caused by constantly carrying a heavy weight on one arm may

FIG. 7.



produce deviation of the spine, and a well-marked dorsal curve.

Also, lateral curvature occurs in rickets. Rickets, however, is a general disease which involves not the bones only, but every tissue of the body. It is sometimes

hereditary ; but much more frequently it is developed after birth, through exposure to cold, and through insufficient or improper food. The whole skeleton may become more or less softened, and the bones curved through mechanical causes. Especially the lower limbs become curved, whereby the pelvis is rendered oblique, and spinal

FIG. 8.



curvature follows. In this form of disease, not only is spinal curvature induced at an earlier age than when it arises from debility, but it is of an infinitely more severe form. In the case from which Fig. 9 was taken, rickets was first developed at eight years of age, and the child was thirteen years old when the drawing was made. Here

will be observed flat-foot, knock-knee, and obliquity of the pelvis, as well as spinal curvature. The lower limbs were first bent; then the pelvis was tilted to one side and became flattened, and subsequently the spine became curved.

She and her sister were patients at St. George's Hospital. They were stunted in growth, and of unhealthy appearance. Until eight years of age they had been as healthy as the rest of the family, when they were seized with this form of disease, which gradually crept on. They were well cared for in every way and were lodged in Kensington Palace, the parents being servants at the Palace. It was therefore somewhat difficult to account for this access of disease; for every surrounding circumstance appeared to be in favour of health. Distortion, whether of the spine or of the limbs, is never so great in childhood from any other cause as when it is induced by rickets.

In a large number of cases it is found that where a child has not been brought up at its mother's breast, there is a tendency to rickets. And where a child has been fed with farinaceous food and at the same time nursed at the breast, the same occurs. These children early develop flat-foot. But if a child is brought up on farinaceous food or on any of the mixtures known as infant's food, rickets more or less is pretty certain to be induced. Enlarged epiphyses are sure to be found; with, perhaps, a rickety cranium and also an anæmic condition, with night sweats and a waxy complexion. Such a child has a soft, yielding fibre, and it becomes flat-footed as soon as it touches the ground and begins to walk. This observation refers to the poor and the rich alike.

This rickety condition of the spine is sometimes, though rarely, developed without the long bones being affected. There will then probably be seen also rickety deformity of the ribs and of the sternum. Rickety limbs are found without a rickety spine; or, on the other hand, the whole skeleton may be affected with rickets. Rickety

flat-foot or a curved tibia or femur will cause the pelvis to become oblique, and the spine to become curved. But a rickety spine alone, and without other portions of the frame being affected, is very rare. As Sir Benjamin Brodie stated, "it was the prevailing opinion formerly, and

FIG. 9.



I believe that some hold the opinion still, that the common cause of a lateral curvature of the spine is a rickety condition of the bones. This view of the pathology of the disease is not confirmed by the specimens preserved in

museums of morbid anatomy, and no one who has seen much of these cases in the living person can doubt that the fact is otherwise. We are not, therefore, justified in regarding rickets as the common, or even as a frequent cause of spinal curvature: nevertheless it is the cause of it in a few instances.”*

Also, there is a form of curvature which is transmitted with a strumous diathesis, and which pervades certain families. There are families of which every member is thus afflicted—some severely, and others slightly; but all, or nearly all, are affected with spinal curvature.

Again, spasmodic action of the muscles of the back, neck, and shoulder, will draw the spinal column away from the mesial line. This is sometimes, though very rarely, found to exist at birth; and occasionally it is met with together with other manifestations of morbid action.

Congenital curvatures are sometimes excessive in degree; so that the trunk may be reduced to two thirds of its normal height. When the muscles of one side of the back are alone affected scoliosis results; but when both sides are affected lordosis is produced.

Lately I saw a child of four months, who had a strongly marked lateral curvature with the convexity towards the left side. It was affirmed that this state existed from birth. When the child cried the hands were clenched, the legs were flexed upon the thighs, and the thighs upon the abdomen, and the feet were extended, while the shoulder and the ilium of the right side were violently approximated.

Fig. 10 was taken from one of the most remarkable cases of spasm that I remember to have witnessed. In this instance torticollis and scoliosis were induced to so great an extent, that it was difficult for the patient to remain upright; and, indeed, this position could only be maintained for a very short period. The sketch gives unfortunately a very inadequate idea of this case.

* Lectures on “Distortion of the Spine not connected with Caries, ‘London Medical Gazette,’ vol. xxxviii.

After some duration of spasmodic action, loss of power, partial or complete, may ensue.

In the anencephalous and the hemicephalous infant every variety of distortion of trunk and limb is to be found. And also somewhat similar distortions are seen with hydro-

FIG. 10.



cephalus and hydrorachis. Sets of muscles only may be retracted so as to produce torticollis, talipes, or strabismus ; or, on the other hand, every limb as well as the trunk may be contorted. These and other forms of muscular retraction exist, both as congenital and as non-congenital distortions ; they are each induced by a like cause, and they differ only in degree and in form.

There are still two other very rare varieties of lateral

curvature to which it is only necessary to allude—viz. curvatures arising from malformations of the vertebræ, and those consequent on disease within the thorax. In the former there is occasionally seen, either a deficient development, or an excess of development in the lateral halves of some of the bodies, and indeed of other portions of the vertebræ. There is a remarkable example of this malformation in the museum at Vienna, which I have examined, and which consists of four half-vertebræ, with their half-arches and processes in excess.* These occasioned curva-

FIG. 11.



ture of the sacrum, slight curvature in the lumbar and the lower dorsal regions, considerable curvature in the middle dorsal region, and also a considerable curve in the upper dorsal region. I have seen a somewhat similar instance

* Vide Rokitansky's 'Pathological Anatomy,' Sydenham Society's edition, vol. iii, p. 228.

of deformity in a child of three months old, and other cases at more advanced ages.

Effusion into the cavity of the thorax, which is followed by collapse of one side, is subsequently compensated by enlargement of the opposite side. After the effused fluid has been removed from the pleural cavity, the ribs sink to the compressed lung, the lung itself being bound down by adventitious membranes, so that it cannot expand. Thus the side becomes flattened and the lung remains contracted. Compensation afterwards follows, through amplification of the sound lung and enlargement of its containing cavity. When curvature arises from this cause the curve occupies the entire dorsal region. Its appearance is shown in Fig. 11.

This curve is formed in the recumbent position ; and it forms as a single curve, and without a compensating curve, until the patient rises from his bed. Then compensating curves are formed, pretty much as in other cases ; and then also rotation of the vertebræ commences. Thus, a lateral curve is first formed, and it is followed by rotation.

CHAPTER II

FORMATION OF SPINAL CURVES

LATERAL curvature may commence either in the lumbar, the dorsal, or the cervical region. No curve, however, can remain single, but secondary curves must form, to restore equilibrium. Thus a lumbar curve will run into a dorsal, a lower dorsal into an upper dorsal, and an upper dorsal into a cervical curve. By observation it may with precision be stated which is the primary curve, and for purposes of treatment this knowledge is of the greatest importance.

When a curve is formed from below upwards, there will be found an oblique pelvis. This will have been caused by some difference in the lower limbs—one being shorter or weaker than the other, and in consequence the spine becomes inclined towards the lower side of the pelvis.

This lumbar curve is ordinarily the primary curve. A primary cervical curve is caused by torticollis, and a primary dorsal curve may be caused by any considerable increase of or difference in the muscular power of an upper extremity, such as follows ankylosis of a shoulder, amputation or paralysis of an upper extremity, or without any marked increase of muscular power, by carrying a heavy weight, such as a child, always on one arm, the spine becomes curved with its convexity towards that side. This dorsal curve terminates above in a cervical curve and below it blends with a lumbar curve. The cervical curve induces an upper dorsal curve. A spinal curve cannot remain single, except in the horizontal posture; but compensating curves are always formed to restore equilibrium. A primary curve is never entirely compensated

by a secondary curve; and therefore other curves form above and below it. Thus, a dorsal curve is always followed by one above and another below it; and a lumbar curve not only occasions a dorsal but also a sacral curve, and a cervical curve is compensated by the weight of the head, which is inclined towards the opposite side, and by an upper dorsal curve. One curve runs into the other, so that as one forms another is forming.

Thus a primary curve may be lumbar, dorsal or cervical, and the position of the primary curve is determined by the exciting cause of distortion; while compensation takes place through the formation of secondary curves, which are produced by the muscular effort to maintain the erect position. These curves are always undergoing change and becoming more rigid and compressed, until at length the height of the trunk is much diminished.

In the commencement of spinal curvature the length of the lumbar curve depends in great measure on the tilting of the pelvis. When this is much the curve is short, and when it is slight the curve is longer; so that as obliquity of the pelvis increases the lumbar convexity becomes more prominent, and the curve itself represents the segment of a smaller circle. As rotation increases the curves become compressed, and the whole spine may lose several inches in height.


Perhaps the modes in which spinal curves are formed will be more readily understood by means of the following diagrams (Figs. 12 and 13).

Suppose the lines A, B, C, D to represent the pelvis and the spinal column in their normal relations; namely, the spinal column forming a right angle together with the pelvis; and let E F represent the oblique pelvis, and G B the inclined column still forming a right angle with its base. How is equilibrium to be maintained? It is obvious that the line G B, if continued, would render equilibrium impossible; and, therefore, a series of curves about the perpendicular line A B are formed to maintain equilibrium.

It will be seen, however, in Fig. 13, that the cervical curve corresponds to the line G B. Such, indeed, must be more or less the case until equilibrium is restored through increase of the spinal curves. When such increase has taken place, as is shown in Fig. 14, then the cervical curve approaches the perpendicular line.

The symptoms and the external characters of spinal curvature vary according to the region in which the primary curve is formed.

FIG. 12.



When, from any affection of the lower limbs, the pelvis becomes oblique, the spine must be curved in the lumbar region. The ilium is depressed on the side of the lumbar convexity, and the loin falls in on the opposite side, through the recession of the lumbar vertebræ in the formation of the lumbar curve. From the concurrence of these two conditions the hip on the side of the lumbar concavity becomes prominent. This appearance is constant, and it is one of the most striking symptoms which may

be observed : it may always be seen even before a secondary dorsal curve is fully formed. This is shown in fig. 15.

Again, when curvature commences in the dorsal region, or when a compensatory curve has been formed in this region, the shoulders are no longer placed on the same level: one is raised while the other is depressed. And the scapula becomes unduly prominent on the convexity of

FIG. 13.



the dorsal curve: the shoulder is popularly said to be "growing out." This increased prominence of the shoulder is due, in part, to the increased angularity of the ribs, and partly also to muscular action.

In the formation of a lateral curve the vertebræ become twisted, or rotated; and consequently the angles of

the ribs on the convexity of the curve project abnormally, while the ribs themselves are rendered more horizontal in their direction and are more widely separated from each other than in their normal condition ; but on the side of the concavity the ribs become oblique and are depressed, so as to lie one upon another, or even to overlap one another.

FIG. 14.



There is considerable change also on the anterior surface of the thorax. On the side corresponding to the concavity of the spinal curve the ribs become unduly prominent at the lower part of the thorax, while they are still more prominent on the dorsal aspect on the side of the convexity. Hence the greatest diameter of the chest, in an advanced state of dorsal curvature, is in an oblique direction from behind forwards.

The muscles on both sides of the spine are somewhat pallid, and wasted. In the commencement of curvature they are tense on the side of the convexity, and later, as rotation increases, they become prominent through the projection of the transverse processes. At length they become atrophic through disuse. In the concavity

FIG. 15.



of the curve they are, for the most part, weak, relaxed, and inactive.

It has been shown that at birth the physiological spinal curves do not exist, but that they are due to the erect position ; and that thinning of the intervertebral cartilages is due to unequal pressure. This compression is such that a man of middle stature, who has maintained the upright

position during the day, is found to have lost nearly one inch in height. Such is the case with the normal—antero-posterior—curves. And, when lateral curvature takes place, a somewhat similar course is observed. A slight lateral curve is first formed: it is not yet persistent however, but curvature is effaced after the horizontal position has been maintained for some hours; just as in the antero-posterior curves the height is restored after the recumbent position has been observed for some hours. At length, however, the intervertebral cartilages which enter into the curve remain more or less compressed, and the curve continues to be formed even in the horizontal position.

During the incipient stage of curvature, then, the intervertebral cartilages become compressed laterally, and they recover their form when the superincumbent pressure is removed. When, however, this compression is continued from day to day, the intervertebral substances lose in a measure their elasticity, and do not recover their full form during the period of repose, but remain somewhat compressed and wedge-shaped.

In the act of walking there is a lateral undulating movement of the spine from the pelvis to the head, which corresponds to the alternate acts of raising the feet from the ground. As the weight of the body rests on the right foot, for instance, the pelvis is raised on the same side, and the undulation commences in the spine with the convexity towards the left side, and there is a compensating undulation in the dorsal region towards the right side. Again, as the left leg is brought forward, the pelvis becomes horizontal, and as the weight is transferred to the foot so the pelvis is raised on the same side and rendered oblique, and the lumbar curve forms to the opposite side, as has been already explained.

Together with this undulatory motion of the entire column there is also a certain slight lateral movement of each vertebra upon the intervertebral substance, especially in the lumbar region of the spine. This is a movement of rotation in the vertical axis of the column, and it varies

from right to left with each step.* When however, the column is no longer erect, but inclines laterally, this normal movement of the vertebræ is checked. Then, compression on the side of the concavity commences to be permanent. The immediate effect of this compression is seen in a certain amount of stiffness, or rigidity of the spine, which shows inability to rotate towards the concavity, and consequently for the bodies of the vertebræ to replace themselves exactly in the central position.

Confirmed lateral curvature is not purely a lateral deviation of the spine, except when it takes place after pleurisy with exudation, or after abscess within the thorax during recumbency. Under all other circumstances the vertebræ, which are involved in the curve, become rotated upon their axes and displaced laterally as the curve increases. After pleurisy or abscess, and during the formation of the thoracic curve, the patient will have been entirely recumbent. It is only when he rises from his bed that compensating curves are formed, and that rotation of the vertebræ takes place.

The bodies of the vertebræ are not all in the same measure rotated; but those are most rotated which are nearest to the centre of the curve, and that vertebra which is central is the most rotated and the most wedge-shaped. This is shown in Fig. 19, where the vertebra in the centre of the lumbar curve is represented as wedge-shaped and rotated to the extent of one fourth of a circle, while those above and below are both less rotated and less wedge-shaped. And in the same manner the intervertebral substances which enter into the curve are reduced in thickness.

In rotation the bodies of the vertebræ are turned towards the convexity of the curve, and the spinous processes towards the concavity.

In a severe form of curvature the bodies of the verte-

* These movements may be studied with advantage in young acrobats, in whom exaggerated positions of the trunk and limbs will display these movements of the vertebræ.

bræ undergo such rotation as to acquire a lateral instead of their normal direction. But, although the bodies of the vertebræ are rotated, the spinous processes undergo comparatively slight change in direction, and even it may be so slight as scarcely to indicate a lateral curve. This is shown in Figs. 16 and 17, especially in the dorsal region. Here the spinous processes scarcely indicate a lateral curve although the bodies of the vertebræ are much rotated.

Perhaps the course of the spinous processes in the lumbar portion of the spine, as represented in Figs. 18 and 19, is even more remarkable than in the dorsal region considering the greater rotation which has taken place.

Thus, it is shown that a spiral twist of the spinal column may exist to a very great extent, through rotation of the bodies of the vertebræ, without the apices of the spinous processes describing a corresponding curve.

This question of rotation has attracted much attention ; and every one who has touched on the subject has propounded his own solution of it, with more or less confidence. Many have theorised on the subject, and opinions differ as to the cause of rotation and the time of its commencement. Rotation occurs both in primary and in secondary curves, but it is always considerably more advanced in the primary than in a secondary curve.

It has been stated by Dr Dods,* who was one of the first to draw attention to the subject, that lateral curvature always commences with rotation of the bodies of the vertebræ, and that it is produced by muscular action. Mr Barwell seems to have accepted this view.† Others have thought that a difference in the density of the anterior and the posterior portions of the vertebræ would account for rotation. Mr Alexander Shaw believes that certain changes in the articulating processes are due to unequal pressure occasioned by rotation. He says : " In lateral curvature of the spine we have a distinct demonstration that the articulating processes give way more extensively

* 'Pathological Observations on the Rotated or Contorted Spine, p. 30.

† 'On Curvatures of the Spine,' 3rd ed., p. 27.

FIG. 16.



FIG. 17.



FIG. 18.





FIG. 19.



than any of the other parts of the column. This is evinced by the rotation which the spine makes in its perpendicular axis, at the same time that it inclines laterally.”*

Mr William Adams adopts Mr Shaw’s view, and says, “I believe rotation to precede any lateral deviation of the spinal column, and its cause and mode of production to have been accurately explained by Mr Alexander Shaw.”†

Dr Judson has lately expressed his views of rotation, and his explanation is so ingenious that I shall refer to it at some length. Also, he illustrates his theory by an instrument, which will probably be familiar to many in this metropolis, as it was lately extensively exhibited by Dr Sayre. Dr Judson writes as follows: “In the present unsettled state of the question I propose an explanation as follows:

“The rotation of the vertebra in lateral curvature of the spine consists in an unequal lateral displacement of the body and the spinous process, the former being free from lateral attachments and therefore departing from the median plane, while the latter is held in the median plane by its muscular and fibrous attachments.

“While the spinous processes of the vertebral column are held in the median plane of the trunk by the strong and active fibrous and muscular structures which are attached, on one hand to the posterior portion of the spinal column, and on the other hand, to the shoulders, the thoracic and abdominal parietes, and the hips, the anterior portion of the vertebral column is comparatively free from fibrous and muscular attachments.

“The bodies of the vertebræ have motion upon each other by the intervention of the intervertebral substance. They are unlimited in lateral motion, except by the intervertebral substance, the anterior and posterior common ligaments, their slight visceral and costal connections, and from being connected by their pedicles with the posterior portion of the vertebral column, the component parts of

* ‘Medico-Chirurgical Transactions,’ vol. xvii.

† ‘On Curvature of the Spine,’ 2nd ed., p. 150.

which are bound together by ligamentous and muscular structures.

“When the median plane of the trunk curves to the right or the left under the vertical pressure which is the direct cause of lateral curvature, their freedom from control allows the bodies of the vertebræ to fall away from the median plane, to the right or the left, while the posterior portions of the column are held in the median plane by their muscular and fibrous attachments. This deportment of the two components of a vertebra, its anterior and posterior portions, is rotation in the vertebra as a whole.

“The distinguishing feature of the explanation of rotation here proposed is the recognition of the fact, heretofore overlooked, so far as I am aware, that the posterior portion of the vertebral column, being a part of the dorsal parietes of the chest and abdomen, is confined in the median plane of the trunk, while the anterior portion of the column, projecting into the thoracic and abdominal cavities, and devoid of lateral attachments, is at liberty to, and physiologically does, move to the right and left of the median plane.”

Dr Judson concludes his paper thus :

“The immediate cause of the curvature is an inability to support the superincumbent weight. What tissues are at fault, and why they fail at a certain point of development, and in females more than in males, are questions which are not included in the subject of this paper.”*

From this paper we are to understand that the spine yields from above downwards, in the same manner as the curves are produced in the instrument which is figured by Dr Judson on the first page of his pamphlet, and that it yields from “inability to support the superincumbent weight,” and “curves to the right or the left under the vertical pressure which is the direct cause of lateral curvature.”

This instrument is doubtless based on experiments made by Professor Hermann Meyer, in which he showed that

* ‘The Cause of Rotation in Lateral Curvature of the Spine.’

by vertical pressure on the spine of a seven months' foetus, he could produce lateral curvature with rotation. And at fourteen years of age the same state was produced, but not to the same extent, until the anterior longitudinal ligament was removed: after the removal rotation was complete. But at twenty years of age rotation could not be produced even after removing the anterior common ligament.*

Dr Judson was doubtless familiar with this paper of Prof. Meyer's; and it will be observed how closely his opinions agree with those of Meyer. It will not be difficult to show that an error underlies the theory; and that when reduced to practice it does not work.

As a rule a lumbar curve is first formed through obliquity of the pelvis, and the pelvis is rendered oblique through some inequality of the lower limbs, either in length or strength. In the early days of a flat-foot or a knock-knee, there will be observed a tendency for the pelvis to fall to the same side, and also a deflection of the lower portion of the spine with its convexity to the same side. When, however, the patient is placed in a recumbent position, so that the superincumbent weight is removed, the flat-foot and the knock-knee and the oblique pelvis and the deflected spine disappear: the arch of the foot is restored, the knee no longer deviates from the normal line, the pelvis becomes horizontal, and the spine occupies the median line. It need scarcely be said that at this period rotation has not commenced. When rotation commences the spine does not resume immediately its normal position.

A left lumbar curve, then, depends on a faulty condition of the left lower extremity, while a faulty right extremity gives rise to a right lumbar curve. And of the upper curves each one follows a regular course, in the same manner as the lumbar curves. Thus spinal curves are formed with precision; so that the original defect being given it is easy to describe the curves which must follow: curves do not form to the right side or to the left as though by accident, but almost with mathematical precision.

* "Die Mechanik der Skoliose," Virchow's 'Archiv,' vol. xxxv, p. 225.

So soon as spinal curvature commences the axis of the trunk is changed, and the column is no longer poised in the vertical line on the heads of the thigh bones. The superincumbent weight being no longer transmitted in the normal axis of the trunk, but falling on the side of the concavity of the primary curve, unequal pressure causes first compression of the intervertebral substances which enter into the curve. After a variable time this compressed condition of the intervertebral substance continues more or less, and is not removed by ordinary recumbency. The body of the vertebra then begins to move laterally, through the pressure on the inner, or concave side of the curve, and in consequence of pressure being removed from the convexity. And through this compression on the side of the concavity, a gliding, or rotating movement of the bodies with the intervertebral substances is established. This compression acts first on the intervertebral substance, and secondly on the body of the vertebra; and causes thinning and lateral displacement, or rotation.

Rotation then, is immediately due to abnormally increased pressure on the concave side of the spinal curve, and to removal of pressure from the convexity.

Rotation can never be a primary movement, but must always be a sequence of lateral deviation.

It is essential to rotation that the spinal column shall be bent.

The position which is assumed in commencing curvature is that of stooping forward with the head inclined towards the side of the lumbar concavity, and subsequently, as equilibrium is restored, to the side of the convexity.

In health, the column is in itself perfect and able to sustain the weight which is superimposed in ordinary conditions; but when its base is rendered oblique, the column yields laterally: it is no longer perpendicular, and is therefore unable to support the superincumbent weight. Thus a lumbar curve forms, the centre of gravity is shifted, and the superincumbent weight is transmitted

to one side of the median line, rather than through the normal axis of the body.

Then, doubtless, the curved spine is unable to bear the superincumbent weight. It curves through obliquity of the pelvis, and not through vertical pressure, and it is unable to support the superincumbent weight because it is curved.

Curvature and rotation commence from below upwards in 95 per cent. of cases of lateral curvature, and not from above downwards, as is represented by Meyer and Judson.

The experiments to which I have alluded show the immense importance of the anterior common ligament. They also demonstrate that curvature precedes rotation, and that rotation can only arise out of curvature.

When curvature commences from above its progress is slower than when it commences from below. Yet the curves form in regular sequence, and the stability of the column is impaired; but it is never so much weakened as when its base has lost its horizontal direction and has become oblique.

In figs. 18 and 19 it will be seen how the bodies of the vertebræ are rotated towards the convexity of the curve, and that they have become wedge-shaped. That vertebra which is the most central is also the most compressed: it may lose more than half of its thickness. It will further be seen that the spinous processes are somewhat curved towards the concavities of the curves, but not in the same degree as the bodies are rotated into the convexities. They may even scarcely be removed from the middle line of the trunk, notwithstanding that the bodies themselves are much rotated.

Therefore, in defining a spinal curve, we must not be guided alone by the line of the spinous processes; but the condition of the entire trunk must be considered. To consider the spinous processes alone would mislead.

Figs. 20 and 21 represent an articulated spine, where the anterior and the posterior surfaces agree perfectly, and where the vertebræ are piled one above another with great

care after the fancy of the constructor, but unfortunately without the slightest regard for truth. Such is the manner in which articulated spines are frequently put together ; and it tends to perpetuate an error. It is only fair to say, however, that the vertebræ having been taken asunder and the intervertebral cartilages removed, it is no ordinary

FIG. 20.




FIG. 21.



puzzle to put the several parts together again as they had grown.

The transverse processes on the convexity of the curve, by rotation of the vertebræ, become prominent and cause the erector spinæ to project ; while, on the opposite side, they can be only imperfectly felt. Also, through the pres-


sure caused by rotation the articulating processes are partially absorbed.

The ribs necessarily follow the altered positions of the vertebræ to which they are attached, and undergo a movement of rotation backwards on the convex side of the curve, so that their angles are rendered more prominent, and they become more horizontal in their direction, while the intercostal spaces become wider than in their normal state; but on the concave side the ribs sink and become flattened, the intercostal spaces also become more or less effaced through overlapping of the ribs, and the ribs themselves are carried forward and become prominent on the anterior and lower part of the chest. On both sides of the chest the ribs are flattened. Also, on the convex side of the curve, in consequence of the rotation of the bodies of the vertebræ into the convexity and the flattening of the ribs, the lung is compressed. Through these changes in the form of the thorax and others which are coincident with them, the capacity of the chest is diminished. Further, the appearance of distortion is much increased by the prominence of the scapula. On the convex side of the curve this bone is thrust up, and is placed obliquely, through the increased angularity of the ribs; and it is still further raised by muscular action. It is also occasionally observed that the edge of the latissimus dorsi does not pass over the lower angle of the scapula, but that it lies below it, the scapula being raised.

In an ordinary case of lateral curvature the pelvis is oblique. The ilium on the side of the lumbar concavity is raised and carried backward, and on the opposite side it is depressed and carried forward. When the superincumbent weight is unequally transmitted to the ground, the pelvis becomes slightly flattened (fig. 22). And, when affected with rickets, the pelvis becomes flattened from above downwards, both through the superincumbent weight and by the resistance of the lower limbs; so that the space between the promontory of the sacrum and the symphysis of the pubes is diminished.

When distortion of the spine is considerable, and of long duration, some flattening of the pelvis ensues in consequence of the unequal transmission of the superin-

FIG. 22.



cumbent weight. Thus, the ischium and the os pubis become somewhat flattened on that side on which the weight of the body is most thrown, namely, on the side of the convexity of the lumbar curve.

When difference in length of the extremities has given rise to curvature, such difference as is occasioned, for instance, by a rickety femur or tibia, by genu valgum, or other such cause, flattening may take place on the side of the shorter limb, that is to say, on the side of the convexity of the lumbar curve. The flattening which occurs, except in a rickety pelvis, is slight.

In addition to cervical, dorsal, and lumbar curves, the sacrum may be curved in a sense the reverse of the lumbar curve; and the ilium on the side of the sacral concavity is then carried forward. Also, through these changes, together with the torsion of the spine which has occurred, the promontory of the sacrum encroaches slightly on the pelvic cavity.

Distortion of the spine is never stationary, but it advances slowly, until, at length, ankylosis takes place between the bodies of the vertebræ, and also between the

articulating processes. Ankylosis may ensue after absorption of the fibro-cartilages, or it may occur through deposition of bone along the bodies of the vertebræ, in riband-like masses or in the form of nodosities—stud-like masses, connecting two or more vertebræ, without the previous removal of the intervertebral cartilages.

The anterior surface of the thorax presents changes which are scarcely less striking than those which have already been mentioned. Thus, on the side of the convexity, the ribs are flattened and depressed; and, together with them, the position of the mammary gland is changed: its position varies, being depressed from a quarter of an inch to an inch or more. This necessarily depends both on the situation of the curve and on its extent. The gland retains, if not entirely, almost its normal relation to the ribs, and is depressed together with them.

On the side of the concavity of the thoracic curve, the ribs and cartilages become prominent; so that the projection is not unfrequently spoken of as a tumour. Sometimes pain is felt in this position; and not unfrequently this is the only seat of pain in scoliosis. This prominence is occasioned through the rotation of the vertebræ and the consequent thrusting forward of the ribs and their cartilages.

The sternum also deviates somewhat from its normal position, its lower extremity being directed towards the side of the convexity of the thoracic curve.

The viscera, both of the thorax and the abdomen, are compressed, and their functions become deranged in every severe form of lateral curvature. Such is the case, indeed, before distortion reaches its most severe stage. But the thoracic viscera suffer especially, being surrounded by bony walls. On examining the skeleton, one is apt to wonder how the functions of respiration and circulation can be carried on, so great is the change from the normal condition.

Rachitic scoliosis seldom appears until disease is fully established in the long bones, and in the pelvis. The

nature of the affection can, therefore, rarely be doubtful. A rachitic tibia or femur, by inducing obliquity of the pelvis, may cause spinal curvature. This is an every-day occurrence. The curve is not, however, therefore rachitic. Rachitic scoliosis and ordinary scoliosis have little in common but the existence of a curve and the compression of viscera consequent upon distortion.

Rachitic scoliosis commences in the bony structures. First, the periosteum becomes unusually vascular, tumid, and adherent to the bone. The structure of the bone itself is then somewhat swollen, and its vessels are filled with blood. The cells, too, of the spongy parts of the bones, and of the epiphyses of the long bones, become dilated and filled with sanguinolent fluid. Eventually the cell-walls are broken down, and the medullary cavities become chambers filled with this fatty, sanguinolent, dark-coloured matter. This fluid is found both in the cavities of the bones and between the periosteum and the bone; and at length the bones become soft and cartilaginous, being, in part, deprived of their earthy constituents, so that they are easily cut with a knife, or as readily yield to pressure or to muscular force, and become misshapen. Wherever pressure is greatest there is the most distortion. So in the spine it is found that the vertebræ which are most implicated in the curve are not only much compressed, but they are twisted on their axes, and through pressure become misshapen.


Through the weight of the trunk also the pelvis is flattened and narrowed in its conjugate diameter; and this tendency to flattening is increased by the existence of lordosis, which, as a rule, exists together with scoliosis in rickets (Fig. 23). The ribs on both sides of the thorax are flattened, and they are distorted into every conceivable shape.

To this softened and tumefied condition another even more important yet succeeds. The bony substance becomes more and more firm, the outer surface being hypertrophied and of ivory hardness; so that the distorted

limb or spine, however misshapen, must so remain : change is impossible.

In no other form of spinal curvature does the pelvis become so distorted as in that which results from rickets.

FIG. 23.



A case of this description is on record where a ball, one inch in diameter, could not pass through the brim of the pelvis. The Cæsarean section was performed by Dr Hull.

In the second place I will proceed to examine the result of these changes on the thoracic viscera, as well as those which take place in the structures which are attached to the trunk itself.

The thoracic space, on the convex side of the curve, is diminished by the flattening of the ribs, and by the rotation of the bodies of the vertebræ ; and the heart is consequently somewhat displaced towards the concave side of the curve. Respiration may be considerably affected ; and through imperfect expansion of the chest and lungs, the right side of the heart becomes dilated, and the blood is insufficiently aërated.

The aorta follows the inflections of the vertebræ, being bound down to the spine by its branches ; and thus it follows the curves of the spine. Its course under these

circumstances, is well shown in a preparation numbered 3416 in the museum of the Royal College of Surgeons. In a practical point of view, this course of the aorta may appear to be a matter of only small importance. It deserves to be remembered, however, for in an emaciated person, with the convexity of the lumbar curve towards the right side, the aorta may be felt immediately beneath the finger, lying out of its normal course, and on the right side of the umbilicus. I have known ideas of aneurism arise in connection with severe spinal curvature with rotation. I was lately consulted in such a case, where it was supposed that a very severe dorsal curvature to the right side was complicated with aneurism of the aorta. It occurred in a gentleman, aged sixty-five.

Together with severe lumbar curvature, there is always found obliquity of the pelvis. This is not a simple tilting to one side, but there is at the same time a slight movement of rotation of the pelvis itself, which, indeed, is necessitated by the circumstance of the lumbar curve and the rotation of the lumbar vertebræ; so that the anterior superior spinous process of the ilium is not only raised above that of the opposite side, but it is also thrown back; while on the opposite side it is depressed and carried forward. In the female this obliquity is of less importance than in the male. In the male, however, the triangular ligament of the urethra, together with the rest of the pelvis, is twisted, and consequently the direct course of the urethra behind the ligament no longer corresponds with that in front of it. This twisted condition of the urethra may cause a serious impediment to the introduction of a catheter into the bladder, even where stricture does not exist. Should stricture of the urethra be superadded it would be important to remember the oblique position of the pelvis. I have known more than one instance where, without the knowledge of this fact, it would have been very difficult to introduce a catheter into the bladder. Whenever in cases of severe lumbar curvature it is necessary to use an elastic catheter, this is always withdrawn

moulded into a sigmoid curve, similar to the urethral curve.

In the case to which allusion has been made above this difficulty occurred. It was found impossible to introduce a catheter until this twisted condition of the urethra was explained and overcome.

CHAPTER III

TREATMENT

BESIDES all those measures which are usually taken to improve or restore health, it has to be considered what special means are adapted to the removal of lateral curvature.

The chief causes of lateral curvature of the spine are, as I have endeavoured to show, those which act on the pelvis through the lower limbs, and in all such cases the lumbar is the primary curve. The first endeavour must then be to restore to the pelvis its horizontal position. This may be done in the recumbent position; when the flat-foot or the knock-knee, &c., which have occasioned obliquity of the pelvis may be then treated. But the recumbent position is not absolutely necessary, for the causes of obliquity of the pelvis may be removed equally well, though the time occupied will be longer, without recumbency. Until the trochanters are brought exactly to the same level it is of little use to attempt to act on the pelvis and the lumbar curve; and until this state is produced walking should be prohibited.

I do not propose to enter upon the question of constitutional treatment for debility, nor for rickets, nor will I touch further upon the treatment of such conditions of health as are induced by lengthened residence in climates unsuitable to the race; which by forcing growth, and advancing the period of puberty, diminish strength and unfit such young people to associate on equal terms with those who have never been subjected to such influences. But I may call attention to the necessity of guarding

against inordinate exercise, long walks, and much standing. Such girls need reclining chairs and suitable sofas.

In examining for spinal curvature the shoulders and the hips must be uncovered, and the whole length of the spine should be traced with the eye, as well in the stooping as in the erect position. Stooping may bring into view a rotated spine, which a practised eye even shall not discover in the erect position ; and, also, the loin may then be seen to have fallen in, whilst in the erect position there shall be no apparent defect.

If the lumbar curve is so slight that it can be removed in the recumbent position, spinal supports are not necessary ; but gymnastic exercises, to bring the muscles of the trunk and extremities into action, will be sufficient to restore the spine to the median plane of the trunk. These exercises alternating with rest, together with such constitutional measures as every one understands for invigorating the general health, will suffice to remove incipient curvature. But, although this statement is made in the fullest confidence of the efficiency of such treatment, it should at the same time be said that such cases need the most watchful care, lest they should pass on as they easily may to the next stage, namely, of rotation. When rotation has taken place recumbency and gymnastic exercises will no longer avail to restore the spine to the median line of the trunk ; but mechanical support is then indicated, together with other measures which will now be mentioned.

The treatment of spinal curvature should commence so soon as the curve begins to be formed ; or as Sir Benjamin Brodie very justly said "The treatment of the disease cannot be begun too soon after the first signs of spinal curvature are perceptible."* For if curvature is allowed to progress deformity must go on increasing until equilibrium is restored by the formation of compensating curves. However slight the curve may be, it demands

* Lectures on "Distortion of the Spine not connected with Caries."
'London Med. Gazette,' 1846.

immediate attention. It is supposed by some that, as if by magic, a child may grow out of lateral curvature. No hope can be more delusive. When the column inclines only slightly to one side, it requires all the care that can be bestowed, so that curvature shall not increase.

If it be a fact that one curve is first formed, and that others are subsequently formed as compensatory of this primary curve—and no one can doubt it who has watched these cases attentively—then it should follow that treatment must in the first instance be directed especially to the removal of the primary curve; for to endeavour to remove a secondary curve without giving efficient support to the primary curve is the most certain mode that could be devised of increasing the original curve. Having determined then which is the primary curve, pressure should be made in that direction which shall tend to restore the positions of the ribs, and also of those vertebræ which have undergone some rotation. This is most effectively done by applying the force to be used below the greatest convexity of each curve, whether a primary or a secondary curve.

The manufacture of a spinal support and its application are matters of considerable importance in relation to curative measures. A spinal support must be so made as to be perfectly fitting in all its parts; it must be light, and so tempered that it shall not bend to the weight of the body, but be bendable with a mechanical wrench. The band about the pelvis should fit exactly, and therefore it must be fashioned to the pelvis itself; and the lateral plates which are carried on levers and moved by compound movements (lateral and antero-posterior), should correspond exactly to the surfaces to be acted on.

That exactitude may be obtained, a cast of the back must be taken, and a skilful mechanist must construct the support. The lateral plates should be so placed that they will apply if possible to the lower portion of each curve and they will be broad in proportion to the curves to be acted on. Sometimes it is not possible to act effectively

in the first instance on the lumbar curve. The treatment may then commence by acting on the pelvis itself and on the lower dorsal curve, with probably a third plate to act on an upper dorsal curve.

The dorsal curves will be acted on through the ribs. The plates therefore should be moulded exactly to the shapes which these have assumed, and they will from time to time need to be remodelled. The force will be applied in the reverse direction to that in which deviation has occurred, and the pressure should, as much as possible, be made to follow the course of deviation of the parts themselves; and, consequently, it should never be directly lateral.

It is necessary that the plates of the support shall be accurately in contact with the curves at all times when the patient is not recumbent, and that as the curves alter the plates shall be changed, both in shape and in position. When the combined forces of a well-adapted instrument are made to act in the directions now indicated—namely, obliquely towards the centre, they tend to unbend the curves. The movement which is thus commenced in the primary curve is often greatly assisted by muscular action; and in this manner the several curves are at the same time or in succession acted on and unfolded.

The action of the spinal support should be that of uplifting, and therefore pressure will not be made on the greatest convexity of the curve, but on the extremes of the arc. By making pressure upwards and forwards and laterally, rotation may be overcome and the curves themselves opened. This action induces extension of the spine; and there will be found an immediate increase in height to the extent of three fourths of an inch to one inch, and even more.

It is rare that a cervical curve requires special treatment unless there is contraction of the sterno-mastoid. It is then a primary curve and requires for its removal the section of both the sternal and the clavicular portions of the muscle. Then the head should be supported as well as the dorsal compensating curve. When the cervical curve

has to be acted on directly, whether as a primary or as a secondary curve, pressure will be applied to that side of the head which corresponds to the concavity of the spinal curve. The twisting or rotation which the head undergoes in such cases corresponds to that of the pelvis. When the anterior superior spinous process of the ilium is carried forward, and there exists also a cervical curve, the ear of the same side is raised and carried forward, while the other is lower and carried backward. Thus, with a cervical curve, even without retraction of the sternomastoid muscle, the head is twisted and the face is placed obliquely.


When the principles of construction are understood, the only remaining difficulty is to find an instrument-maker capable of carrying them out. An instrument that goes by the same name may be so constructed as to be useless, or it may be so powerful that there may be even more power than is necessary. First, it must be so framed and fitted that it cannot tilt, and upon this framework move levers with plates to act on the extremities of the arcs. Or a single lever moves on its axis on a central band with pads operating simultaneously in opposite directions, or with a shoulder hoop and a pad to effect the same purpose. Each case demands an instrument exactly suitable to its peculiarities: each one differs, and no two are alike.

When the lumbar is not a primary curve, but follows and is caused by a dorsal curve it is of much less importance than when it is formed as a primary curve. As secondary curves both the lumbar and the cervical require comparatively small attention. There is then little or no obliquity of the pelvis and it is only after considerable deformity has taken place that there is any rotation of the pelvis. In treating the dorsal curve the plate of the spinal support will be applied in the same manner as has been already described. Probably the lumbar curve will not require to be supported, but the same object may be attained by applying a plate to the pelvis itself, where it is

worn with much ease ; and again a third plate will be applied to the upper extremity of the dorsal arc.

As examples, Figs. 24 and 25 represent casts, taken in September and in the following January ; the first before treatment commenced, and the second when it was nearly complete. It will be observed how great is the rotation both in the lumbar and in the dorsal region in Fig. 24,

FIG. 24.



and how great is the deformity. Before this patient came under my care she was placed in a Sayre's jacket. This she wore for five months, during which time it was once re-applied. When it was applied the curvature was very slight, but on removing it at the end of five months the deformity was such as is represented. Then she was

immediately brought up to me.* At the expiration of five months, namely, in January, the spine was not only nearly straight, but it was mobile.

I am unable to understand how any one can pretend to treat cases of lateral curvature of the spine who holds the opinions expressed by Mr Adams, namely, that "where lateral curvature of the spine exists in any marked degree, and before it amounts to an external deformity, it is essen-

FIG. 25.



tially an incurable affection by any and every method of treatment."† For, unless after rotation has taken place the spine can be restored to the median plane of the trunk, such cases are unworthy of the attention of any

* I shall be happy to show these casts and the patient to any one who is interested in this subject.

† The 'Lancet,' 1866, vol. i, p. 95.

surgeon. But rotation, according to Mr Adams, takes place at the commencement of curvature, and even before there is any lateral deviation of the column. He says, "I believe rotation to precede any lateral deviation of the spinal column."* It may well be asked then, for what purpose does he propose treatment; and of what does his treatment consist?

Friction may be employed with great advantage, the patient lying on the face, and on a flat, resilient surface. It should be made with the open palms along both sides of the spine; thus acting on the entire loin and the dorsal region. This friction should be made by two strong arms, with all the force that can be borne, especially on the side of the convexity; and some lubricating substance may be employed to save the integument from injury.

Whilst curvature is being removed, attention should be directed to the development of muscular power by means of exercises; such as swinging, riding, swimming, and the like. These alternating with rest, will, if judiciously employed, rapidly produce a marked increase of muscular power, as well as improvement in the general condition of the patient. By swinging is meant suspension by the hands. It should be continued only so long as neither to induce fatigue nor pain.

A spinal chair should be used during the day; and the spinal support will then be worn both with comfort and advantage during the whole day. The inclination of the chair should be such as can be borne for a considerable period with ease; and it should never be such as to cause inconvenience, otherwise a lateral inclination of the trunk will be speedily assumed to overcome this inconvenience. Thus, the chair may be raised or lowered as circumstances require; and in it all studies may be pursued.

A sloping seat, such as is recommended by Mr Barwell, tends to increase pressure on the side of the concavity of the curve. Mr Barwell calls it "a device for lifting that side of the pelvis which is abnormally depressed." In

* *Op. cit.*, p. 150.

sitting, except where distortion is very great, and then of the pelvis itself, the pelvis is not oblique ; but in standing only. But if the pelvis were from any cause oblique, a cushion, as is recommended by Dr Little, might be employed.* The cause of obliquity in spinal curvature is not to be found in the pelvis, but below the pelvis, and therefore in sitting the pelvis for the most part becomes horizontal. Mr Barwell gives an illustration of the effect of the raised seat on the curved spine.† It is impossible that such a condition as is there depicted, *i. e.* the curves absolutely and neatly reversed, could be produced by any seat in any length of time. A curve might be produced easily enough in a healthy spine, by such means.‡ But, by raising the pelvis on the side of the convexity of a lumbar curve, the trunk is necessarily tilted over to the opposite side, and pressure is thereby increased on the side of the concavity ; where, indeed, the intervertebral substances are already compressed. The illustration shows the spinal curves exactly reversed. But to avoid exaggeration, Mr Barwell uses the phrase “the spine *tends* to assume a contrary curve.” “Therefore,” he continues, “I must not be supposed to say that an established lumbar curve can be at once inverted by such means, or even that it will be immediately effaced.” My contention is that such means do not tend to diminish a spinal curve ; but that by raising the pelvis beyond the horizontal line on the side of the convexity, not only the lumbar curve, but also the compensating curves must be increased.

For these reasons the spinal chair in ordinary use, the back of which can be raised or lowered at pleasure, is not only pleasanter to use, but more effective as a means of treatment than a tilted stool.

* ‘On Spinal Weakness and Spinal Curvatures,’ p. 72.

† Op. cit., p. 126.

‡ Prof. Bauer, in his suggestive work, “Lectures on Orthopædic Surgery,” states his belief “That a disturbance of the centre of gravity *per se* is not sufficient to cause permanent deviation of the spine,” but that superadded there must be the peculiar condition predisposing to scoliosis, p. 150.

With regard to the employment of bandages in the treatment of spinal curvature I would express the opinion that they are useless for cure; and unfortunately they are not harmless. They cannot be kept applied; and they are to be found where they are not intended to be. But, even if they could be fixed they would be useless after rotation has taken place. And with regard to the plaster of Paris jacket recommended by Dr Sayre, I have had opportunities of seeing it applied by himself, and of watching the effect. In every instance where the jacket was used, the curve was increased and the height of the trunk was diminished. But in some instances, after its application, there is complete loss of power in the trunk and the lower extremities, so that the patients can neither stand nor sit as a result of the treatment. One who is again under my care had become so powerless in consequence of this treatment, that she could not move in bed nor raise her limbs from the bed. She is a young lady, 23 years of age, who having worn a spinal support for some months was nearly cured; but her friends becoming impatient and thinking that something more might be done for her, consulted a hospital surgeon who is in the habit of applying these jackets. He applied the jacket and it was worn for five months. After that time a poro-plastic splint was applied for a twelvemonth. Then he found himself compelled to give up the case. I was again asked to visit her and found her in bed and unable to move. She had lost all power in the trunk and extremities before the jacket was removed. Unfortunately these sad results are not infrequent. At the present moment I have two such cases under care. They are both very bad cases: one is of sixteen months' standing, and the other of twenty months. In both there is complete paralysis of the lower limbs and of the trunk. In each case the jacket was applied by hospital surgeons. I do not desire to enter minutely into the consideration of the cases which I saw with Dr Sayre; but in all my experience of treatment of spinal curvature and spinal disease I have never known

results in the least approaching such misfortunes. In one case, that of a strong young man with angular curvature, sloughs formed, which were immediately followed by hæmaturia and pneumonia; and in a case of lateral curvature, the loss of height and increase of deformity made it positively necessary to discontinue the treatment. These cases were both under Dr Sayre's own care. But, besides these, I am almost ashamed to allude to dozens of cases which I have been compelled to see, which have been subjected to this treatment. The jacket has been applied to infants of tender age, of some few months or years, and sometimes it has been left unchanged for six months and more. Such was the case in a child of six years from Cornwall, and again in another of ten months from Clifton. In the latter case the jacket was applied when the child was three months old, and it was not removed for seven months. I saw the child wearing the jacket which it had worn for seven months. It nearly died before it could be relieved of its bonds. A more hopeless little cripple could scarcely have been manufactured. The jacket is a very dirty appliance, and not only is it unsuitable and injurious, but it is a shameful abuse of what is termed treatment.

Every back splint, under whatever name, is simply useless in lateral curvature. No one supposes that a Sayre's jacket, or a poro-plastic splint, or a splint constructed of any other material can prevent rotation of the vertebræ and the increase of lateral curvature. Instead of being cured those who submit to such treatment soon find to their sorrow that they become every day more and more crooked. Even if exercise were not permitted rotation would increase, but with exercise it proceeds more rapidly than if no apparatus were worn. The worst cases of lateral curvature that have come under my care during the last few years, have been treated with some sort of splint before I have seen the patient. I entirely agree with Dr Bauer, who says, "All the braces with which I am familiar seem to be useless, for

they all act upon the ribs, and upon these alone. It strikes me that it would be much more easy to bend and permanently deform the ribs than to effect the reduction of the spine by these means" "I cannot conceive how the gutta-percha splint of Schildbach, or the "elastic muscles" of Davies, could make any difference in the result."*

Whilst I write a lady brought her daughter to me with the following history :

This young lady, aged twenty, was under the professional care of a well-known physician and of an equally well-known surgeon, who had treated her for seven years for various ailments. The surgeon sent her to an instrument maker for an appliance for a malady quite unconnected with the spine. But the instrument maker, doubting the surgeon's diagnosis, requested that he might see her back. He immediately found a lateral curve of the spine, and recommended a poro-plastic splint. The physician and the surgeon were communicated with, and instantly agreed to this recommendation.

What I complain of is this: *neither of them have ever examined this patient's spine.* But, notwithstanding, they immediately approved of the application of the splint. On examining her spine I found that there was no curve, and that the entire spine was in a perfectly normal condition. Many of these splints are applied in sheer ignorance, and because there is some pain perhaps in the back.

There are circumstances connected with this case which oblige me to protest strongly against the indiscriminate use of back splints in lateral curvature of the spine. If one had been applied in this instance a series of lateral curves would have been manufactured as in the case mentioned at p. 54 and figured 24. I have now had an extensive experience of these manufactured deformities, and I hope to be excused in bringing the subject before the profession. Spinal curvature has immensely increased, and especially severe cases have increased, since back

* Op. cit., p. 178.

splints and bandages have been in vogue. It is equally unjustifiable not to treat a spinal curve when it is forming, and to encase a back in a splint where a curve does not exist, thus to increase or create a spinal curve. Splints give no lateral support and cannot therefore prevent rotation, but on the contrary they invariably increase rotation, and consequently increase lateral curvature and diminish the height of the trunk. I possess carefully taken notes of forty cases where measurements were kept, and where increase of deformity was proved. And it is not alone increase of deformity that takes place, but it is such rapid increase as could not take place unless a splint or a bandage were worn. Without giving lateral support they cause wasting of the muscles of the back, and they increase the undulations of the body in the lumbar region, and therefore produce increased rotation of the vertebræ.

Again, numberless cases are not treated because the parents are advised that a child will "grow out" of a spinal curvature. In this matter members of the profession have something to blame themselves for. It is not difficult to understand that with a slight curve some doubt might arise as to the absolute necessity for interference, but when curvature has already become severe, hope is still expressed that gymnastics and time will together perform a cure. The evil day is only postponed. Such patients are certain to return worse than when first seen.

Curvature increases greatly through neglect, especially the upper dorsal curve becomes prominent and angular, the scapula being raised on the projecting ribs; while on the side of the concavity the ribs fall away, being flattened. Thus, one shoulder is much raised above the other, giving the appearance as of a hump on the shoulder, with the face carried to the opposite side.

I should gladly have left this part of the subject at this point, but I feel bound to notice a lecture on "Lateral Curvature of the Spine;* and as it was spoken *ex cathedrâ*, I hope to be excused if I refer to it. I will not follow

* 'British Medical Journal,' May 25th, 1878.

the lecturer in his theory of rotation, for this may be dismissed in as few words as Mr Fisher has bestowed on it.* He proceeded to describe the treatment to which he was about to submit a patient, and spoke thus: "Now, I will proceed to raise the left foot so as to bring the pelvis straight, and you see at once a diminution in the lumbar curve, which of course cannot be rectified in a day; still less can the dorsal curve and rotation be put right without assistance, and that I propose to give by applying a plaster-of-Paris jacket after Sayre's method." And he continued: "Now, why should the dorsal curve be almost invariably to the right and the lumbar to the left, and not the converse? The reason is, no doubt, that the majority of mankind are right-handed, and that the right limb being more developed is heavier and the muscles which support it stronger than on the left side." Perhaps he would be surprised to hear that a left-handed person has the dorsal curve to the right exactly in the same manner as the right-handed person. Yet such is the case. And he added, "But in addition to this we must take into account the peculiar type of respiration in the female," and he developed a theory upon "the important respiratory muscle, the serratus magnus;" the whole of which theory, if not wholly a fallacy, is a somewhat doubtful subject for a lecture-room. Then "spinal supports" were discussed, of which he said, "the principle of all is the same." "Theoretically," he considered the principle "is very satisfactory," and even it was "conceivable that if a spinal support could be made to grasp the patient it might do some good." But he contended that they "slip up and down" as the levers are moved.

I would suggest to the lecturer to change his instrument maker. Certainly, with such an instrument as is described, no good can be done. But to return to the patient: she was fixed up, and a bandage was applied, neither the "dinner pad," nor the "cotton wool over the breast," nor another very important piece of the apparatus being

* Ibid., June 1st, 1878.

forgotten, namely, "strips of tin on the projecting side." Then this young woman was dismissed, and the lecture was concluded thus: "In a fortnight's time, gentlemen, this patient will come before you again, and we shall then be able to judge of the amount of improvement which is likely to take place."

A second lecture has not been published, or probably we should have been told what changes had taken place.

It will be observed that strips of tin were placed on the projecting side. Let us suppose that they were placed on the opposite side also. Then a bandage was rolled firmly round the body. Thus the pressure was at least as great on the side of the concavity as on the convexity. But, I would ask, what more is needed to increase rotation? As might be expected, this is the result of the application of the bandage.

There is nothing absolutely new in the mode of extension as it is applied by Dr Sayre. The same apparatus, with a slight difference, is figured and highly recommended by Mr Stafford.* He called it the "spine elongator." But it dates much farther back than Stafford. Glisson introduced this practice in the seventeenth century.† And the book in which he introduced the practice of swinging in spinal curvature is yet much appreciated. In Germany "Glisson's swing" is known and used, and it has been in use ever since Glisson's time.

Extension alone and without other means is useless in

* 'On the Diseases of the Spine,' p. 80.

† "*Artificialis corporis suspensio* perficitur ope instrumenti ejusdem pensilis ex faseis ea arte formati, ut pectus sub axillis complectatur, caputque sub mento alia faseia circumdet, manusque binis ansis exequat, unde corporis onus partim à pueri manibus, partim à capite, partim ab axillis in aëre pendulum sustineatur ita, ut ab adstantibus non sine voluptate huc illuc impellatur. Etenim ad ossa curva restituenda, ad articulos deflexos erigendos ad eurtam corporis staturam elongandam, conducit. . . . Aliqui, ut magis distendantur partes, caleos plumbeos pedibus adaptant, et contractiori corporis lateri pondus adjiciunt, quo facilius partes ad æqualem longitudinem extendantur. Hoc autem exercitium solis robustioribus competit."—'De Rachitide,' p. 415, 1671.

lateral curvature of the spine. Dr Sayre has such merit as attaches to the practice of using swathing bands in addition to this extension. And having applied the plaster-of-Paris jacket he insists that "the patient, whilst wearing the plaster-of-Paris jacket, should practise self-suspension every day once or twice, and from three to five minutes at a time."

The effect of the suspension is, of course, to loosen the bandage, so that in three days the knitted jersey can readily be drawn out and replaced. There can, therefore, be no adequate support to the spine, and nothing to prevent increase of rotation. The exercise of the muscles is praiseworthy, and has been practised certainly for two centuries in this country. The objectionable part is the bandage; it causes wasting of the muscles of the back and those of respiration, and it increases deformity.

There never has been devised since the world began, such an admirable arrangement for the production and the increase of spinal curvature.

Mr Reeves says, "The plaster-of-Paris bandage is worse than useless in any form of lateral curvature."† Where the bandage is left on for a fortnight there is a considerable increase in the curve, and the curve increases in proportion to the time that the bandage is worn. This curve, as shown in Fig. 24, was almost nothing when the bandage was applied. This was continued for the space of five months, at the end of which time no one could desire to see a worse curvature. So long as the pathology and treatment of lateral curvature are taught as in the lecture to which I have referred, so long must these cases multiply and be severe as they have been during the last three years.

I have not seen any treatment that will compare favorably, even remotely, with the mechanical treatment which I have advocated. This can never occasion such results as those above alluded to; but the beneficial results may

* 'Spinal Disease and Spinal Curvature,' p. 101.

† The 'Lancet,' November 15, 1879.

be seen every day, as in such cases as those recorded on p. 54, and shown in Figs. 24 and 25.

Thus, whether treatment is pursued in the upright or in the recumbent position, or in both, the aim should be to unbend the curve. This point has been illustrated by Jules Guérin* and by my late colleague, Mr Lonsdale.† The question is how best to carry it out? Extension is quite useless. This was long since proved by that most able teacher, Guérin; who, by his report to the Government in 1848, to the effect that longitudinal extension alone must be inoperative, stopped a mode of treatment, which if not quackery was nothing.

Pressure must be so applied as to act on the rotated bodies of the vertebræ and on the extremes of the arc, to unbend the curve and to remove pressure from the side of the concavity. Lonsdale carried this out in the recumbent position, but he lost what he had gained when the patient again resumed the spinal support, which was a modification of Tavernier's; for it was unequal to continue the action.

By placing the plates of the spinal instrument as I have indicated, the same action is continued in the upright and in the recumbent position, and treatment may be continued almost as well in the upright position as when the patient is recumbent. Thus the extremes of the arc will be acted on, whether the curve be formed in the cervical, the dorsal, or the lumbar region, by acting on the head, the shoulder, the lower part of the thorax, or the pelvis, as is rendered necessary by the nature of the curves and the cause of deformity.

* 'Mémoire sur l'Extension sigmoïde et la Flexion dans le Traitement des Déviations latérales de l'Épine.'

† 'Observations on the Treatment of Lateral Curvature of the Spine.'

CHAPTER IV

ANTERIOR CURVATURE OF THE SPINE

ANTERIOR curvature affects, for the most part, the lumbar region of the spine, and is an abnormal increase of the physiological curve of that region. It occurs also, however, in the cervical region in infants. It is then due to rickets and to deficient muscular power, and sometimes also it is due to caries. When anterior curvature takes place in the lumbar region it is, for the most part, of rachitic origin. Sometimes it is hereditary, and then, also, it depends on a rickety condition; and, again, it results from congenital dislocation of the heads of the thigh-bones.


Anterior curvature is for the most part limited to the lumbar region of the spine, the dorsal and the sacral portions being implicated only inasmuch that their normal curves become exaggerated. Occasionally, however, a part, or even the whole, of the dorsal curve may be reversed; and then the anterior curve will consist of the lumbar vertebræ together with the dorsal.

In the lumbar region this distortion may vary from a slight increase of the normal curve of this region to a development which is a source of great weakness. Through increase of this curve the obliquity of the pelvis is augmented, the anterior wall being carried backwards and the sacrum and coccyx being raised, so that their concavities present more directly downwards, and the superior portion of the pelvis inclines forwards. The sacro-lumbar articulation, in consequence, is no longer found immediately above the interfemoral line, but in advance of that line. Thus equilibrium is disturbed, and it becomes necessary,

for its restoration, that the weight of the upper part of the trunk shall be thrown behind the vertical line of the body. In this manner a compensatory antero-posterior dorsal curve is produced.

The external appearances presented by anterior curvature then, are such as are indicated by the changes in

FIG. 26.



the skeleton above mentioned. They are represented in fig. 26. The lumbar region is rendered remarkably hollow, while the dorsal is rounded; the nates are raised, and the head is thrown back; the abdomen is unusually prominent, and the stature is necessarily stunted.

Causes.—As a congenital affection it is combined with

monstrosity, with spina bifida, with rickets, and with dislocations of the femur.

An hereditary disposition to lordosis is sometimes observed; one or more members of a family being affected. It may be the only sign of a rachitic tendency, or a more severe and general form of rickets may present itself. In some members the cachexia is so marked that the whole skeleton partakes of it, and no portion of it remains unaffected. When the pelvis is much contracted and anterior spinal curvature is at the same time considerably developed, natural parturition may be prevented, and the Cæsarean section, or other such means, may become necessary for the removal of the foetus. Such instances are recorded by Ramsbotham, Kilian, and others. In these cases softening of the bone had allowed excessive deformity of the pelvis to be produced.

Whatever tends to increase the obliquity of the pelvis will give rise to this form of curvature. Thus, it is a necessary consequence of congenital dislocation of the heads of the thigh-bones (upwards and outwards), Fig. 27. A child was lately under my care for congenital dislocation of the thigh-bones, in whom lordosis was very marked. Indeed, so much was this the case that the congenital dislocation had been entirely overlooked, and attention had been directed to the spinal curvature alone—a result only of the dislocation.

The child from whom the drawing was taken, had been kept constantly recumbent for three years before I saw her, and had been treated for spinal curvature alone.

Lordosis may be induced equally by unreduced non-congenital dislocation of the thigh-bones. Thus it may follow rheumatic inflammation and subsequent dislocation of the head of the thigh-bone on to the dorsum ilii. In these cases as in those last mentioned, great efforts are made to sustain equilibrium; the extensor muscles of the trunk are rendered tense and prominent, and, standing out firmly, they leave, especially towards the lower part of the spine, a deep sulcus in the median line.

FIG. 27.



Some years ago I saw, together with Mr. Brookes at Shaldon, in Devonshire, a similar deformity, but arising from a totally different cause, and I allude to the case because it is rare. Fatty degeneration of the muscles of the abdomen had taken place, and the abdominal viscera hung suspended consequently in a huge, tumid paunch; the weight of which had, in part, occasioned the distortion in question. This distortion was one of the most remarkable that I ever witnessed. The effect was increased, also, by fatty degeneration of many other muscles, such as the trapezii, rhomboidei, serrati, latissimus dorsi, the pectoral muscles, and those of the lower extremities. Many of these muscles were so far destroyed that they could not even be traced. In this instance the unsupported pendulous belly, together with the loss of muscular power, seemed to give rise to lordosis, in a somewhat similar manner to that in which the gravid uterus is sometimes known to act.

Bearing heavy weights around the neck and shoulders will also induce this form of curve. That a considerable burden may thus be slung, borne upon the neck and shoulders and suspended in front, it is necessary that the shoulders shall be thrown back, and that, consequently, the loins shall be projected forwards. If this practice is continued, the increased lumbar curve becomes more or less permanent. I have also known prolonged gymnastic exercises produce a somewhat similar result.

The pathological changes which are induced are chiefly as follows. The lumbar spines are approximated; the articulating processes of the vertebræ are pressed forcibly into contact one with another; the posterior ligaments become somewhat shortened, and the ligamenta subflava lose something of their elasticity; the bodies of the vertebræ undergo absorption and the intervertebral substances are compressed posteriorly; while expansion of these substances takes place on the anterior surface in the convexity of the curve. The extensor muscles of the trunk are strongly developed. Anchylosis of the articular processes

of the vertebræ, with or without fusion of the bodies, may at length take place, through which the distortion is rendered permanent.

Anterior curvature is, for the most part, limited to the lumbar region of the spine; the dorsal and the sacral regions being implicated only so far that their normal curves become exaggerated. Occasionally, however, some few of the lower dorsal vertebræ may become involved together with the lumbar curve; or, indeed, the entire dorsal curve may be reversed, and the anterior curvature will then consist of the lumbar vertebræ together with the dorsal. This is a very rare form of distortion. I lately saw an instance in which the sacrum was almost horizontal, and the head was forcibly thrown back, while the anterior curve was formed by the whole of the vertebræ, lumbar, dorsal, and cervical.

Treatment.—It has already been said that lordosis is seldom a primary affection, but that it is the result of disease, or that it arises from the position of parts which are entirely independent of the lumbar region itself. This affection is for the most part developed in childhood. But in childhood the normal curves of the spine disappear in the horizontal position. Hence it is obvious that the recumbent posture must be in itself a powerful agent in the treatment of this form of curvature. Doubtless, whatever the cause of anterior curvature may be, the treatment should be followed whilst the child is recumbent. The supine position is the best in which to place the child; for the shoulders can then be raised, whilst, in addition, the thighs may be flexed upon the pelvis. In this manner an anterior spinal curve may at length be removed.

When this form of spinal curvature has been induced by rickets, it is especially important to observe the recumbent posture; for the upright position tends to increase the spinal curve, and will probably also occasion some distortion of the lower limbs.

Cases occur, however, in which it is not practicable to

observe constantly the recumbent position, and it then becomes necessary to substitute for it a portable instrument, which, receiving the weight of the head and shoulders, transmits it to the pelvis. The spinal column is thus relieved, and the extensor muscles are less violently thrown into action. It is a very imperfect substitute, however, for the recumbent position.

The influence of the muscles in distorting the arms and forearms is well known. Exercises should on this account be forbidden in rickets. Passive movements of the limbs, however, when they are practised gently, are beneficial, both locally and generally; the muscles then tend to increase in size, and the circulation is accelerated.

The treatment of that form of lordosis which is induced by congenital dislocation of the femur is simply the treatment of the dislocation itself. And, those consecutive curves which tend to restore and which help to maintain the equilibrium of the body, which are consequent on Potts' disease and on ankylosis of the hip, for instance, require little more care than is afforded by a well-adjusted support.

CHAPTER V

POSTERIOR CURVATURE OF THE SPINE

POSTERIOR curvature of the spine is a much more common affection than that last mentioned. It occurs in childhood, in youth, and in old age; and is thus one of the most common forms of spinal curvature. In infancy, the cervical, the dorsal, and the lumbar vertebræ are engaged in this curve; while in youth the middle and the upper portions of the dorsal spine are especially implicated; but in old age the spine is most bowed in the upper dorsal region. In infancy, the muscular system is mainly affected; while in later life the skeleton itself undergoes change, and the intervertebral substances become compressed.

Causes.—The causes of posterior curvature of the spine are debility—whether in infancy, in youth, or in old age—rickets, muscular rheumatism, and partial paralysis. Also, some occupations are apt to induce a stoop in those engaged in them. Thus it is, for instance, with watchmakers, engravers, embroiderers, writers, shoemakers, and others, whose occupations require a stooping position. This position, which is at first irksome, becomes easy, and at length more or less permanent. Those also who are subject to asthma acquire a stoop, which even becomes diagnostic of the disease.

Debility, especially in infancy, is the commonest cause of posterior curvature. The curve, for the most part, occupies the entire length of the spine, from the occiput to the sacrum, the spine projecting backwards. The normal curves of the spine have not as yet been formed, for the muscles of the back have not the power to support the trunk, and the head in consequence falls forward. This bowed con-

dition of the spine increases, and the dorsal vertebræ become more prominent; for, notwithstanding its weakness, the child is seldom kept recumbent. Nutrition is imperfect, and symptoms of rickets perhaps begin to show themselves; such as frequent diarrhoea, perspiration about the head and neck, general wasting, and a blanched condition of the skin. The child is uneasy and fretful. Swelling of the ends of the long bones may now be observed; especially of the carpal extremity of the radius and of the tarsal end of the tibia; and the vertebræ will probably assume a somewhat more prominent appearance.

Some of the worst forms of this disease which I have seen have been produced in Russia. Bad ventilation and intense cold are among the causes of rickets, both in the palace and in the cottage in that country. In our own country, and especially in the manufacturing districts, insufficient and ill-assorted food with stimulants, together with damp and dark dwellings, often underground, produce a fearful amount of rickets, which might doubtless be prevented by better sanitary conditions.

The following figure (28) was taken from a Russian boy, who had been born and bred in a palace in St. Petersburg, the son of a great noble. It is perhaps the worst form of posterior curvature that I have seen.

In youth posterior curvature occurs not unfrequently, during rapid growth, as a consequence of debility. At this period the lower cervical and the upper dorsal vertebræ are especially affected. A stoop which is thus acquired, either during rapid growth or after illness, may continue permanently in manhood. Also, those who are near sighted, and who do not habitually wear glasses, are apt to accommodate their figures to their imperfect vision, and may thus acquire a stooping position.

Posterior curvature is not attended with tenderness in the course of the spine, so that firm pressure may be made along the spine without causing pain. Should, however, tenderness be found on examining the spine, destructive

disease, or caries must be suspected. And, indeed, caries is often so insidious in its advances, that irremediable mischief may already have taken place before disease is recognised.

In old age an abnormal increase of the dorsal curve occurs through debility. A similar condition is fre-

FIG. 28.




quently observed, especially among the labouring rural population, consequent on chronic rheumatism.

The pathological appearances differ according to the age at which posterior curvature is observed. In infancy the muscular system is chiefly affected; while in the adult the intervertebral substances become compressed anteriorly;

and in old age fusion of the bodies of the vertebræ, or more partial ossification, may be found. Together with these changes, the whole trunk is more or less affected: the chin rests on the sternum, the ribs are approximated, and respiration is laboured in consequence of diminished

FIG. 29.



mobility. Fig. 29 shows the condition of the thorax and spine in the adult.

The treatment of this affection in childhood is simply to maintain the horizontal position until debility is so far removed that the child has acquired the power to assume a sitting or a standing posture.

A slight form of support enables children to assume a sitting posture. When, however, the head droops, recumbency ought to be maintained; otherwise an apparatus to support the head becomes necessary. Such children should be treated for debility in the horizontal position.

As strength is gained, means may be taken to increase muscular power by well-directed exercises. These may be used either in the horizontal or in the upright position. And as muscular power becomes developed, so the antero-posterior curves of the spine become formed, and the erect position becomes possible. In more advanced age mechanical support is necessary to redress or to support simply the bent spine.

Hereditary cyphosis will seldom be removed by any means: it is, for the most part, conjoined with thoracic disease.

Perhaps the best form of support is that which is made with a stiffened pad, which has been accurately moulded to the lower portion of the curve. This is carried on a framework which supports the shoulders. And by means of rack and pinion joints attached to a band around the pelvis, levers act so as to produce moderate pressure.

DISEASE OF THE SPINE

CHAPTER VI

DISEASE OF THE SPINE AND ANGULAR CURVATURE

DISEASE of the spine occurs very frequently among children, following upon a slight accident, and it occurs less frequently among adults. No period of life is exempt from this disease, but it is essentially a disease of childhood. It is seen more frequently in the male than in the female. Thus of 500 cases which I have tabulated, 297 were males. It may occur in every portion of the spinal column, from the head to the sacrum, but it is more frequently observed in the dorsal than in the cervical or the lumbar region; first, because the dorsal is the most exposed to injury, and secondly, because it is the most extended portion of the spine. Yet it occurs very frequently in the cervical region. In the dorsal region it is attended with the greatest deformity, but in the cervical with more danger to life.

Although an accident usually precedes the development of spinal disease, it would appear occasionally to arise out of a deposit of tubercle in the cancellous structure of the bone, which softens and breaks down. Whether, however, traumatic inflammation precedes the deposit of tubercle, which is highly probable, or not, inflammation when it becomes chronic assumes a strumous character.

In its mode of progress this form of inflammation does not materially differ from that which is observed in other articulations. It is first developed in the body of the vertebra. There are many examples, however, to show

that the inter-vertebral substance may become diseased before the bone.

In an early stage of disease one spot at the seat of injury is painful; sometimes it is acutely painful. This pain in the first instance probably depends on laceration of ligamentous and other fibres. Presently it subsides somewhat, the child cries itself to sleep, and the adult perhaps remounts his horse. Then, especially in the child, it may be forgotten, or, rather, it may not have been realised where was the seat of pain or the nature of it. Thus, in this early stage, disease of the spine is frequently overlooked. The pain to which it gives rise sometimes is styled neuralgia, and it is often supposed to be connected with thoracic disease. Pain may continue to be felt by the child for months, and by the adult for years without the disease being recognised. It is experienced long before any irregularity of the spinous processes is observable, and it is attributed to anything but the right cause. It is very common to see cases where disease has advanced to produce considerable deformity, and where caries of the spine is not even suspected.

The following are cases in point:

CASE 1.—I saw with my colleague, Mr Tatum, Mr H—, who at nineteen years of age was a crack billiard player, and, being unable to ride and to indulge in field sports, devoted much time to billiards and much to painting. At this age he could not rise to straighten himself from the table after delivering his stroke, and would remain bent and with his back almost at the same angle during the entire game, and thus for hours in succession. His back became more rigid, and in the course of years he lost flesh. He saw physicians and surgeons without end, and was supposed to be suffering from rheumatism. He was heir to an entailed estate, and he became possessed of it when he was twenty-seven years of age. He laid out all his available money in setting this place in order, that he might enjoy it, and pass it on to his son. When he was thirty-two it was thought that he had

thoracic disease, and he was sent to Madeira for the winter. There he became worse, and could not bear any movement; so he was slung in a hammock, and in this manner was brought to his house in London. Until this time his spine had never been examined. I saw him some few days after his arrival. He was sitting in bed with his shoulders and head slung forward for support, and I found the dorsal vertebræ carious and extremely painful to the touch, and the patient frightfully emaciated. There was no sign of abscess. Some few days later I was hastily called to him in the early morning, and found him unconscious and moribund, one gallon of pus having escaped from the posterior mediastinum, per anum. As he left no son his estate passed to his cousin, and he left his wife a pauper.

CASE 2.—I saw, with Dr Beale and with the patient's father, a medical practitioner at Liverpool, and subsequently with Dr Little, a gentleman twenty-six years of age, of a strumous diathesis, who two years previously had hurt his back while stooping. His back remained "stiff" and painful from that time. We found the four lower dorsal vertebræ carious, and an abscess in the lumbar region. He had just taken his passage to China, having no notion that he was suffering from disease. He was placed on a couch, and remained on it for three years, until firm ankylosis had taken place. Pain subsided, and his general health improved in a short time after he began to lie down. In the course of ten months the abscess was absorbed. At length he was upright and strong, and tolerably active, with scarcely any projection of the vertebræ. In 1869 he married. Two years later lithotomy was performed by Sir Henry Thompson, and a considerable calculus of phosphate of lime was removed.

After the first few days pain is less acute. The muscles of the back become rigid when the dorsal vertebræ have been injured, and especially those of the neck become rigid when the injury is of that region. Pain continues as long as there is inflammation, and when

it is diminishing it may always be excited by pressure or by percussion. The child moves about carefully, placing his hands on pieces of furniture as he passes through a room, resting with his hands or his elbows on a table, or resting one or both hands on his thighs in walking.

But, also, this position is adopted for the sake of the support which is afforded by it, and that the superincumbent weight may thus be in some measure removed from the diseased structures. At the same time the muscles of the back are thrown into action to fix as much as possible the affected parts, and thus to prevent movement and the pain which is caused by movement; in the same manner as is observed in all disease of articular surfaces. Muscular rigidity takes place to a much greater extent in the cervical than in the other regions of the spine; and for this reason that in a normal condition the mobility of the cervical portion is greater than that of any other portion of the spine.


At length inflammation passes on to suppuration, the cancellous structure of the bone breaks down, and there is found a prominence of one or more vertebræ. Caries is then established; and, as disorganisation proceeds, a gap of greater or less extent is formed, according to the number of bodies involved in the disease, and the upper portion of the column falls forward. That vertebra whose body has been most excavated becomes the most prominent, and its spinous process forms the apex of the angular projection (Fig. 30).

In this form of disease the anterior portion of the spinal column, namely, the bodies with the intervertebral substances and the ligaments, is alone liable to become carious. The posterior segment, consisting of the spinous, transverse, and oblique processes, with the pedicles and arches, shows but little disposition to fall into a similar state of disease. Cases are recorded, however, and among others there is one related by Dr Buckminster Brown, of Boston, U.S.*

* 'American Journal of the Medical Sciences.'

The deformity which ensues is greatest in the dorsal, and least in the lumbar region. But the greatest deformity from an equal amount of destruction occurs in the upper dorsal region. This is due to the shape of the vertebræ, as well as to the formation of the vertebral

FIG. 30.



curves; for the bodies of the dorsal vertebræ and the intervertebral substances of this region are thicker behind than in front, or the reverse of that which obtains in the cervical and in the lumbar regions, and deformity is necessarily greater in this than in the cervical or in the lumbar

region, the dorsal spine presenting its convexity backwards. It is also due to the weight of the head. But, also, the dorsal spines are long and superficial; and any increase of the curve increases the prominence of the spinous processes. Thus, loss of substance, when to a small extent only, produces more deformity in the dorsal than in either of the other regions, and is immediately followed by an angular appearance of the spine; and when the chasm is deep the muscles of the back are powerless to support the trunk, and it falls forward. In the lumbar region the vertebræ allow of considerable loss of substance with comparatively small deformity, for they present a concavity backwards with short spinous processes which are buried among powerful muscles, while the bodies are broad and deep.

As the substance of one or more of the bodies of the vertebræ is broken down, the upper portion of the column falls forward. This displacement occurs by reason of the chasm which has been produced in the column: the upper portion of the column may be said to be left unsupported, and therefore it falls forward, so that the spine from the seat of disease may become horizontal instead of vertical. This is especially the case when disease occurs in the lower dorsal and the upper lumbar vertebræ, and where destruction is extensive as is portrayed in Fig. 31.

Many will remember the poor little lad who swept for years the crossing at the junction of Bond Street and Grosvenor Street, and whose spine represented exactly such an angle as is represented in Fig. 31. This is the situation of greatest deformity when the chasm is deep; and the next to it is the upper dorsal region, not alone on account of the vertebra prominens, but because of the weight of the head carrying the upper portion of the spine forward.

The amount of deformity depends much on the rapid progress of disease. In the first instance, the muscles are rigid to fix the diseased portions of bone, so that pain which is caused by motion may be avoided. As destruc-

tion proceeds the trunk would become more and more bent; but an effort is made to preserve equilibrium: the head is thrown back, and the spine below the disease is curved forward, while at the seat of disease the back is rendered more prominent. This is the result of the upright position, and of leaving the spine without support.

When ankylosis has taken place in the recumbent

FIG. 31.



This specimen shows what great deformity may take place, not only without loss of life, but without paralysis; and with only such loss of muscular power as is caused by position.

position with only slight projection, a curve will form above as well as below the ankylosis so that equilibrium may be restored.

In the stage of irritation the muscles of the back are rigid and tend to fix the spinal column and prevent motion so long as pain exists; but they undergo atrophy in an advanced stage of disease and when displacement has taken place. Thus, the amount of angular deformity which takes place depends, first, on the extent of destruction of the bodies, and secondly, on the locality

of the disease. But the extent of the disease depends on the treatment to which the patient may have been subjected. It is possible that little or no apparent deformity shall remain, even when the upper dorsal region has been affected. Cyphosis may disappear entirely or almost so in the prone position, and bony union may form, and a strong and upright column result.

This occurred in a notable case, J. N—, aged twenty-six, who for many years had suffered from disease in the lower dorsal region. There was only slight pain on pressure, but deformity was considerable. Four vertebræ were implicated, and there was pus in the right iliac fossa. In the prone position, deformity was eventually entirely removed. Then, a retaining apparatus having been fitted, he turned about as he pleased, but he was always recumbent. The pus was absorbed soon after he began to lie down. The column became strong, and within three years of rising from his couch he took charge of a large parish in the south of England, which he has administered for several years, and where he is a most active incumbent. When I first saw him he was walking daily, and was not wearing any retentive apparatus. He suffered much pain, and always leant on a stick or on the table when he had the opportunity. He had lost three inches in height.

When disease is established in the cervical region, the head is somewhat thrown back, and is more or less fixed; and the muscles are thrown into action to prevent motion and to avoid pain, while the head seems to subside on to the shoulders. This is especially observable in children. It may also be noticed in adults. Also, the muscles of one side of the neck are sometimes more contracted than on the opposite side, and the head is in consequence inclined to one side or the other, as in wry-neck.

Or, on the other hand, the head may be held forward or it may be turned to one side or the other and so be fixed, as in wry-neck. And, indeed, disease in this region may so closely simulate ordinary wry-neck that an error of diagnosis may easily be committed.

Generally, however, the head falls back, and the child is unable to raise it. In these cases, any movements of the head, especially sudden or rotatory movements, cause excruciating pain; and even sudden death has been occasioned by such movements—namely, in cases where the atlas and the axis were involved in the disease. Under such circumstances the spinal cord in its especially vital portion may be crushed between the atlas and the odontoid process.

This traumatic wry-neck occurred in the following case :

Caroline S—, aged forty-three, was admitted into Princess's ward with caries of the fifth cervical vertebra. The head was firmly fixed and inclined to the right side: the ear was drawn towards the right shoulder, and the chin was projected in the opposite direction. The muscles on both sides of the neck were rigid, but they were much more contracted on the right than on the left side. Such, then, was the position of the head when the patient entered the hospital. After the head had been efficiently supported on pillows for some few days, the muscles became relaxed and the wry-neck disappeared; for irritation had ceased.

Occasionally, two portions of the spinal column are simultaneously attacked with caries. Such cases are rare, however, and their treatment differs in nothing from others which are more simple.

In the dorsal region, the great length of the spinous processes, and the forward curve which this portion of the column naturally assumes, tend to make any projection of the dorsal spine remarkable. The thorax, also, undergoes considerable change in its external form: the sternum is rendered prominent, while the ribs are compressed laterally, and project backwards together with the vertebræ. All the viscera, both of the thorax and abdomen, have to accommodate themselves to their altered parietes. The heart and lungs suffer especially: the heart's action is irregular, and respiration is hurried.

When disease occurs in the lumbar region, it is sometimes attended with considerable pain, but rarely with much deformity. Fig. 32 is intended to show disease and disorganisation of a lumbar vertebra, where the column has fallen forward and encroached on the medullary canal, without, however, producing deformity.

FIG. 32.



Abscess, when it occurs, is always a serious and often a fatal complication of spinal disease. Happily, under treatment, disease for the most part terminates in ankylosis, without the formation of abscess. Rather it should be said that pus is formed in small quantities only in connection with the diseased vertebræ, and that it is absorbed into the system as it is formed, without collecting and burrowing, as it must do, when formed in sufficient quantities. The pus lies in contact with the carious bone and occupies the gap in the bodies of the vertebræ (see Fig. 33). According to circumstances the abscess remains

stationary, or the sac enlarges and the abscess passes downwards by the sides of the bodies of the vertebræ, or it passes down on either side of the bodies, the sac having bifurcated. When the abscess is double the two parts seldom present at the same time; but one will probably only appear when the other is already tense. This is especially the case with psoas abscess.

In the neck the pus passes between the muscles and comes to the surface, burrowing among the fasciæ, sometimes below the mastoid process of the temporal bone, more frequently lower down and nearer to the spine; but commonly it gravitates along the bodies of the vertebræ. In the thorax it lies behind the costal pleura, and perforates the diaphragm through absorption, having become adherent to it, and thus it gains the psoas muscle. Here lying behind the peritonæum, a kind of sheath is formed for the abscess by the fascia iliaca. Thus it proceeds downwards to Poupart's ligament. A large accumulation of pus may form in the iliac fossa, or perforating the abdominal walls the pus will pass with the conjoint tendons of the psoas and iliacus muscles and point at their insertion. When it arrives at this spot, at the top of the thigh, it is known as psoas abscess; and it still retains this name whatever its downward direction may be. It may pass in the course of the Sartorius even to the knee; or having emerged with the psoas and iliacus tendons it may be diverted somewhat inwardly, and still more rarely somewhat outwardly.

Instead of passing in the course of the psoas muscle, however, the abscess on passing through the diaphragm may be deflected backwards, and perforating the abdominal parietes will show itself as a broad and low swelling in the loin, on one or on both sides of the vertebral column. It is now known as lumbar abscess. And when disease is situated in the lumbar vertebræ the abscess will pass upwards and backwards, and perforating the quadratus lumborum will appear in the loin to the outer side of the erector spinæ. Abscess in the iliac fossa will not un-

commonly be absorbed, but psoas abscess is much more rarely absorbed. I have known the iliac fossa to be cleared in numerous instances by absorption.

It is occasionally seen that the contents of an abscess come to the surface along an intercostal space. And instances are known where adhesions having formed between the abscess and the lung, the contents of the abscess have been voided by coughing and expectoration. And also, in a similar manner, it may discharge itself through the bowel or through the bladder.


When, unfortunately, pus accumulates in such quantity that an abscess shows itself externally, whether in the groin, the loin, or elsewhere, it should never be allowed to open spontaneously, but the contents of the sac should be evacuated by means of the aspirator; or the opening may be made antiseptically. Then, tension having been relieved by the withdrawal of pus the opening may be closed. This operation may be repeated as it becomes necessary. Without motion, probably the abscess would not have formed. If motion be allowed after the abscess has been emptied it will again fill. It will probably refill after it has been emptied, even though perfect quiet be observed; so that the aspirator may have to be used more than once.

I have only once experienced any difficulty in emptying the sac of the abscess with the aspirator. In this instance the needle became choked, not with flocculi, but with solid matter, after the fluid portion had been drawn off. On withdrawing the needle nearly one pound of solid matter, having the consistence and the colour of putty, was extruded from the wound. It was extruded continuously in coughing. The wound healed, as it usually does when the needle is withdrawn.

When several vertebræ become affected, so that the chasm in the column is considerable, an angle more or less acute is produced, and the canal itself may be encroached upon, as in Fig. 33. This encroachment may occasion pressure on the cord. Should it take place

suddenly or increase rapidly, it will be followed by paralysis. When, however, deformity increases slowly through the resistance of the structures in the posterior segment of the column, the cord becomes so tolerant of pressure that probably loss of power will not ensue: it is protected by the theca vertebralis, and it accommodates itself wonderfully to the pressure to which it is submitted. In these cases partial paralysis is apt to occur when motion is per-

FIG. 33.



mitted. And, again, power will be restored by continued rest of the body. There was lately, in Drummond Ward, an instance in point. In this case disease was limited to the lower dorsal region, and there was very slight displacement; so little, indeed, that when recumbent scarcely any projection of the spinous processes could be observed. It was, however, conspicuous enough when the patient stood up. It was not possible to keep this patient always

recumbent: she would occasionally get out of bed. When this occurred, violent twitchings of the lower limbs followed these efforts, and they lasted, perhaps with acute pain, for twenty-four or more hours, and then ceased, and the limbs again became motionless and flexed upon the trunk. Then, gradually the muscles relaxed and the

FIG. 34.



limbs resumed their normal condition, when motion and sensation were again perfectly restored.

In the case from which Fig. 34 was taken the bodies of at least five or six vertebræ were entirely destroyed; but, notwithstanding the great deformity which was produced, symptoms of compression of the cord were never experienced.

For the most part, when destruction of the bodies of

the vertebræ has taken place rapidly, paraplegia is more or less complete ; especially this is the case if movements of the body are permitted. But when caries has advanced slowly, probably no muscular weakness will result. Whenever motion is allowed in disease of the spine, a sense of weakness or pain is experienced at the seat of disease ; and this is always increased by percussion or by firm pressure. Somewhat later, painful twitchings are produced by movements of the trunk, and afterwards sensation becomes somewhat blunted. Tenderness begins to be felt in the course of the psoas muscle, or otherwise in the loin or in the neck, according to the seat of disease ; and presently a slight swelling, communicating a doughy sensation, may be perceived, which increases according to circumstances to larger dimensions.

When paralysis occurs, motion is first lost, and subsequently sensation ; and it occurs thus for the obvious reason that the anterior columns of the cord, which give off the nerves of motion, are more subjected to pressure and irritation, lying, as they do, more immediately in contact with the carious bone, than the posterior columns, from which the nerves of sensation proceed. Paralysis is not, however, as I have already shown, in proportion to the amount of deformity but rather in proportion to the rapidity of the change which is effected.

Through pressure at the seat of disease the cord becomes disorganised and motion is impaired or lost. Loss of motor power may be complete, and sensation may yet be retained, or, on the other hand, both motion and sensation may be lost. Also, power may or may not be retained over the bladder and rectum. This depends on the amount of disorganisation of the spinal cord, and also on the position of the injury. Again, one extremity may be more completely paralysed than the other. Sensation may be lost as well as motor power ; but it is never lost whilst motor power is retained. When sensation and motion are lost bedsores frequently form, and when the control of the bladder and rectum is lost then it is

difficult to prevent bedsores. During the week whilst I write I have seen the following four illustrative cases.

A young lady of seventeen was suspended and a Sayre's jacket was applied. She lost all power of motion in the lower limbs, and sensation as high up as the ribs, as well as control over the bladder and rectum. Large sores formed over the sacrum and the sacro-iliac synchondroses. The disease was in the upper dorsal region.

A lady, fifty years of age, was thrown out of a carriage two years and a half ago. She was unhurt, and thought no more of the accident. She continued her usual life until within two months of the present time, feeling pains in the back more or less for about two years, but she called it to herself rheumatism, and never spoke of it even to her intimates. Two months ago she was observed to be sitting for a long time in one position and without moving, and it was found that she had lost sensation and power of motion as high as the waist. She retained full power of the bladder and rectum. Large sores formed over the sacrum, and also on each side over the sacro-iliac synchondrosis. There was no deformity of the spine, but tenderness in the lumbar region.

A lady, aged forty-eight, developed carcinoma in the left breast. This was removed. Two years later there was found a tumour involving the uterus and its appendages. Presently after there was entire loss of sensation and motion, and of power over the bladder and rectum. With great difficulty sores were prevented forming over the sacrum. The bodies of three lumbar vertebræ were destroyed, but without deformity.

A lady, aged forty-six, fell, three years ago, and struck the sacrum. She lost all power of motion in the right lower extremity, but could stand firmly for a short time and with a little help on the left leg. Sensation was perfect. She had complete power over both bladder and rectum. There was no deformity of the spine, but there was acute pain on touching lightly the lower dorsal vertebræ. Loss of motor power commenced twenty months after the fall.

These sores occur especially when sensation is lost. There is no pain felt through continued pressure, and therefore there is no inducement for the patient to move. Unless he is repeatedly raised and his position slightly changed, and also unless the skin is kept dry, sores are very rapidly formed. Indeed, it is difficult to keep the surface whole. The back of the heel is also very liable to suffer from pressure, and it is therefore desirable to place the heels in hollow pads.

When sensation as well as voluntary motion is lost spasmodic movements of the limbs are apt to occur. The knees are jerked up, and again the muscles relax, and the limbs are allowed to fall; or, again, the limbs are more violently flexed, and they remain more or less bent until they are replaced by an attendant. These movements are excited by any irritation, such as touching the soles of the feet, any movement of the body, shaking the floor in approaching the bed, or, again, indigestible food, or defecation, or a breath of cold air, or almost anything or nothing will excite these spasmodic movements; and they are extremely disturbing and wearying, and sometimes, though not usually, they excite pain.

At the same time that caries affects the bodies of the vertebræ, adhesive inflammation is set up in the posterior segment of the spine and in the surrounding structures. This tends to consolidate the disjoined vertebræ and to prevent friction. It is seen more extensively in the neck than elsewhere, because there is more motion here than in any other part of the spine, and it is observed least in the lumbar region. In the neck it not unfrequently assumes very considerable proportions. It may become as large as the fist. This formation, by tending to prevent motion, leads to ankylosis; and when consolidation has taken place the boss is absorbed.

Some years ago I saw, with Sir William Fergusson, a young lady who at eighteen years of age had disease of the fourth, fifth, and sixth cervical vertebræ. There was very slight deformity when she was first seen, so that her

friends believed her health would be improved and the spine not further injured by half an hour's walk daily in Hyde Park. She was wheeled to the park, and there took her exercise wearing a support to fix her head and spine. During the remainder of the day she was recumbent. In the course of one month there was developed a boss over these vertebræ as big as a man's fist. From this time onward, and until ankylosis was complete, she remained recumbent, and with the neck so fixed as to be motionless. When ankylosis had taken place the boss was absorbed gradually and very slowly, and in about three years from the period of resuming her active habits it had entirely disappeared. Now, there is very slight shortening and thickening of the neck, but there is scarcely any appearance of remains of the disease, so that even critical observers have failed to detect any abnormal appearance whatever. This lady is very tall, stout, and active. She was married about five years after ankylosis was complete, and she has been married at least ten years.

There is spurious as well as true ankylosis. It consists for the most part of a thin plate of bone or of bone and fibrous matter, which is formed in the course of the posterior surface of the spinal column. It does not fill up the space caused by the removal of the diseased bodies; and being slender it is liable with only slight exertion to be broken through.

Spurious is more frequent than true ankylosis; and when disease is extensive true ankylosis is rare. With spurious ankylosis a certain amount of health and activity may be enjoyed, so that even laborious exercises are undertaken; but there is never such robust health as with true ankylosis. Active exercise, if the spine is unprotected, is liable to terminate suddenly, through the breaking of the band, in paralysis.

It is not uncommon that inflammation shall extend from the spine to the sacro-iliac articulation; or inflammation which assumes a strumous character may be set up

independently of other disease. It may arise from injury or from exposure to cold, and sometimes it results from parturition. For the most part the bones of the pelvis or of the spine are diseased as well as the articulation. I lately saw in the neighbourhood of Chesterfield a case of sacro-iliac disease where necrosis of the right ilium had followed erysipelas. When the disease is localised there is puffiness about the joint with great pain. The pain is much more than that which accompanies spinal disease. Suppuration and necrosis may follow, or ankylosis may take place.

Perhaps the worst form of this disease that I have seen occurred after parturition in a strumous person. The disease after some years was much relieved, and even it was thought to be cured, when it was again excited by fatigue and exposure to cold. This poor lady died worn out with suffering. There was no abscess. I saw her with Dr Hutchinson and Sir William Gull.

In no respect does treatment differ in these cases from that of disease of the spine. This form of inflammation must be regarded as a more severe disease than that of the lumbar vertebræ, and it induces extensive suppuration and hectic, and terminates, in a large proportion of cases, fatally.

The symptoms are recognised with considerable difficulty, so that suspicions of hip disease are apt to cross the mind. And, indeed, it is sometimes difficult to move the hip-joint without exciting pain. Under ordinary circumstances the recumbent position is to be maintained, but resting on the nates is sometimes too painful to be borne. A heavy person may then be slung, and the pressure can in this manner be modified to afford great relief; and still more relief may be given by the additional support afforded by broad linen adhesive bandages or by belladonna plasters applied in broad strips. Suppuration will probably not occur when this treatment is commenced early.

The coccyx is little liable to disease except when induced by injury; but injuries to the sacro-coccygeal joint are common. A fall on the ice, or on to a stair, or other

angular substance, may cause displacement of the bone forwards. It may be displaced at a considerable angle so as to press upon the rectum. There is intense pain after the accident, which is followed by swelling at the joint. Pressure over the articulation is very painful, or rather acute pain is excited by the least touch. Defecation excites pain, and also walking upstairs or walking otherwise than slowly on the level. Running is impossible; and the act of sitting is acutely painful.

All local applications are useless. The only treatment of any avail is to replace the bone. This is effected instantaneously by the finger in the rectum. Usually the bone returns into position with a click. When it has once been replaced there does not appear to be any tendency for it to be again displaced.

At the beginning of 1883 I saw a case where the bone had become displaced, causing great pain, through falling on to the edge of a stair in May, 1881. It was most easily reduced, however, and the pain immediately ceased. But the case that struck me more than any other that I ever saw was that of a young lady, aged twenty-two, who, while skating at Christmas, 1880, fell in a sitting position on the ice. I saw her six months after the accident. She had become pale and thin, and had lost appetite. Lying on the back was painful, and sitting was very painful, so that it was necessary to use an air pillow. Defecation was impossible except with laxatives. She could walk slowly for about ten minutes. Pain extended from the sacro-coccygeal joint, which was highly sensitive, to the first lumbar vertebra. It is rare to see any one so much distressed as this patient was. The coccyx was displaced and pressed on the rectum. On touching the bone and pressing it slightly downwards and backwards it was instantly replaced. At the same instant all pain left her. The effect was magical. In two minutes the expression of her face had changed from deepest despair to joy; and she exclaimed, "I don't know what you have done, but all my pain has gone, and I feel quite right."

It is the only treatment that is necessary, and none other is effective.

The coccyx is said sometimes to be painful in hysterical persons. I have never seen such a case, and I have never heard such pain alluded to except after an accident, nor have I known a case that was not relieved by replacing the bone.

Disease of the first and second cervical vertebræ deserves, perhaps, separate mention, inasmuch as they are shaped differently to all other vertebræ, and they contain within their area a specially vital portion of the medulla spinalis. Fortunately they are less liable to disease than any of the bones of the spinal column. Being protected by the occiput these vertebræ are seldom injured by force directly applied, but being the centre on which the head rotates, the axis, and especially the odontoid process, suffers from violence to the head.

When atlo-axial disease is set up it assumes a chronic form, and softening of the soft structures between the bones takes place, which is followed by ulceration of the cartilages and partial destruction of the vertebræ. Under these circumstances the movements of the head are attended with extreme danger. The weight of the head will cause the atlas to slide forward upon the axis, and the ligaments being ruptured the medulla will be crushed by the posterior arch of the atlas against the odontoid process. Mr Holmes refers to a case of this kind where death occurred instantaneously on the nurse lifting the patient from the bed. It was found post mortem that the transverse ligament was ruptured, and that the odontoid process pressed upon the cord.* But, as in other parts of the column, displacement of bone and consequent compression of the cord may be so gradual that paraplegia may not supervene, and recovery may take place with some slight deformity only. Or, the entire body from the head to the feet may be paralysed, and recovery may take

* 'The Surgical Treatment of the Diseases of Infancy and Childhood,' p. 542.

place. And again, after fracture, with only slight displacement, paralysis may continue after ankylosis has taken place with complete repair, as in Mr Whitmore's case of John Carter, related by Mr Hilton.* In this instance the patient fell from a tree, forty feet, and injured the fifth, sixth, and seventh cervical vertebræ. He was paralysed below his neck, and remained paralysed, although ankylosis of the vertebræ took place with very little deformity and encroachment on the spinal canal. The bodies and arches of the fifth, sixth, and seventh cervical vertebræ became blended together by bone, and the body of the sixth vertebra was displaced. Mr Hilton remarks, "Every one must admire the perfect and level union by new bone which has taken place at the fore part of the spine, and if nature could have been as effective with the spinal cord this patient might have perfectly recovered." It is exactly the circumstance of the sudden violence rather than the gradual compression which constitutes the essential difference in the paralysis of these cases. In either case recovery may take place, but the prognosis is much more favorable where compression has taken place gradually than where paralysis is the result of sudden violence.

An interesting case of displacement in consequence of disease of the first and second cervical vertebræ is recorded by Sir James Paget,† and although there is no history of the case, it is interesting as showing the two vertebræ "firmly united by bone in such a position that the uppermost part of the spinal cord, during and after the process of osseous union, must have been confined within an exceedingly narrow space." It is probable that in this case there was very extensive paralysis.

Also Mr Alexander Shaw's case was one of displacement of the atlas upon the vertebra dentata. Unfortunately, here also there was no history. The preparation showed "the two upper vertebræ and the occipital bone

* 'On Rest and Pain,' p. 102.

† 'Med.-Chir. Trans.,' vol. xxxi.

distorted by disease and united by ankylosis." Mr Shaw states "that all the bones exhibit that strong marking of the processes and eminences which give attachment to the muscles and ligaments, and that solidity of structure which denotes that the individual shortly before death possessed full muscular power." And he infers that ankylosis must have been complete for a long period before the death of the patient, because the ossific matter forming the union had become perfectly dense in texture and its surfaces were rounded and smooth.*

Several years ago a gentleman was thrown on to the crown of his head through his horse falling in the hunting-field. He was supposed to be dead, but consciousness being restored it was found that he was paralysed below the neck. He was brought up from the north of England in the course of a week to be under the care of Sir Benjamin Brodie, and, as he was my cousin, I had frequent opportunities of seeing him. There was fracture through the axis with slight displacement, and it was believed that some other of the cervical vertebræ were fractured. Ankylosis at length took place between the occipital bone, the atlas, and the axis, and the head became immovable and the neck also remained perfectly stiff. He recovered muscular power perfectly, so that he was as strong a rider and walker as before his accident. He was a fearless hunter, and although he was never again in the first flight, as before, yet he was not far behind. These accidents are usually fatal.

Necrosis of portions of the cervical vertebræ is known to occur as a result of syphilis. I saw, with Dr William Ogle, a patient in St George's Hospital who had an ulcerated pharynx, through which passed three bodies of vertebræ. I saw him extract one of these: it was entire. The others were more or less disintegrated. On passing the finger through the aperture the vacant spaces of the third, fourth, and fifth vertebræ were easily recognisable. There was no deformity of the neck.

* Ibid, p. 290.

Portions, also, of the atlas and of the axis have been expelled in other cases in a similar manner. Such an instance is recorded by Mr Hilton of a patient of Mr Babington's,* where the articular surface of the anterior part of the atlas, and also a portion of the odontoid process of the axis were expelled in coughing. In such cases the head becomes fixed. There are many specimens in the London museums which show ankylosis between the occiput and the atlas and axis.

* Op. cit., p. 99.

CHAPTER VII

DIAGNOSIS AND PROGNOSIS

DISEASE of the spine occurs most frequently in childhood, and perhaps the largest number of cases may be seen in the first and second years. There will be observed in the first instance *pain*, secondly *rigidity*, and thirdly *deformity*.

The first period is short, and is soon followed by rigidity, and the acute pain is succeeded by duller, but yet persistent, pain. Or, should the pain not be absolutely persistent, it is excited by gentle percussion or by light pressure. So long as inflammation exists there is pain. When pain endures and the temperature is high the first stage of disease should be suspected.

After the first few days the period of rigidity is established, and the spine loses its natural flexibility. Especially this is observable when the neck is the seat of disease. Then, also, there is increased heat and swelling. Sometimes this swelling takes place to a considerable extent. Coagulable lymph is poured out, especially in the posterior segment, and the column becomes fixed and motionless. Not only is there more swelling in the neck on account of the greater natural flexibility of the part, but this is more apparent than elsewhere; it may extend to three or four inches in diameter, and be proportionately raised. As ankylosis takes place this swelling subsides, and little or no deformity, but some thickening, may alone remain.

It is only necessary to touch lightly, or to make gentle pressure, to ascertain exactly the seat of disease. It is

quite inexcusable to rotate the head violently, to lean heavily on the head, to jump a child down from a chair, or to make it run up and downstairs. All these means are resorted to to form a diagnosis, but they are entirely unnecessary and barbarous. When pain has somewhat subsided such measures will excite a fresh access of inflammation. And not unfrequently chorea and hysteria may thus be excited. I lately saw a young lady, twenty years of age, who had thus been very considerably examined. Choreic movements were excited, and the case was treated as one of chorea. The lumbar disease was overlooked.

Neuralgia of the spine should be recognised and distinguished from the first stage of disease—the only stage with which it can possibly be confounded. Unhappily it is not alone in the spine that neuralgia is mistaken for disease. I have known such an error occur more than once when excision of the knee-joint has been performed, and all the structures within and without the joint have been perfectly healthy. And I suppose every one is familiar with cases where amputation has been performed for pain in a limb (especially a knee or an elbow), and where the pain, having reappeared on the healing of the stump, it has been chased, morsel by morsel, until there was nothing left to amputate. But the pain still continued about the hip or the shoulder as before, only removed a little nearer to the trunk. Equally so in the spine, neuralgia may deceive. When the hand is brushed lightly over the back exquisite pain is experienced in the neuralgic condition, but the same action excites no pain in disease. More or less pressure is necessary to excite pain in the inflammatory stage of spinal disease, but in neuralgia of the spine pressure is borne more easily than a light touch.

Thus, in the inflammatory stage, and before deformity is apparent, light pressure of the finger along the spine will disclose the seat of disease.

There is a certain tenderness in the lower cervical

region, common enough among young women and those who suffer from exhausting discharges. Also, it is not absolutely confined to the sex. It may be accompanied with swelling. Pressure induces uneasiness rather than the pain which accompanies disease, and there is never rigidity of the spine as in cervical disease. It is occasionally mistaken for spinal disease: more than once or twice I have relieved such a patient of her head-rest.

When the stage of deformity arrives, a prominence of one or more spinous processes is formed. Disease is then far advanced, and partial destruction of one or more bodies has taken place.

All will admit that disease ought to be recognised in its first stage: it is then, however, often overlooked. A child is watched assiduously, but disease is not recognised perhaps until a knuckle of bone is prominent. Then there is probably also an abscess to be seen. Nothing, however, has been done to prevent deformity.

The following is quite a common case, and every one will recognise it. I was lately asked to see a child, aged four, who at three years of age fell from his nurse's arms, but was caught before he reached the floor. There was evidently considerable pain, for the child screamed and continued to scream, and the next and the following days he complained of pain, but he never could specify where was the seat of pain. He was examined, and examined continually, but it always remained doubtful which was the seat of pain. At length pain subsided somewhat, but after that fall the movements were never as they had been previously; but they were made with care and deliberation. At length, a dorsal vertebra became prominent. Then, spinal disease was recognised. Before the vertebra became prominent it was admitted that some injury had been incurred, but it was impossible to say what part was injured.

I found the child moving about in a stooping position, often resting his hand on the thigh; with disease of three dorsal vertebræ, and an abscess in the iliac fossa.

Pain is an invariable symptom in the commencement of spinal disease. A child fails to point it out, but it is always acknowledged readily enough when the finger is placed upon the seat of pain. But after early childhood there is never any doubt as to the seat of injury. A man is thrown from his horse, or his horse falls with him, or he remains in the saddle as his horse recovers himself, and the spine is in each case injured. In very few minutes he is perfectly aware how he is injured. But a child is always tumbling about, for the most part with impunity; but now he strikes his back against a stone, and another bumps himself in falling against the edge of a stair. He is hurt at the time, but soon forgets, and perhaps it is never known by those around that he has done more than he does every day.

Every case of spinal disease may probably be traced to some trivial or more serious accident.

Major S—, aged forty-five, was riding a young unbroken horse over a rough piece of country in New Zealand when they both came to grief. He fell heavily and struck his back. After some little time he was able to drag himself home. In the course of some weeks he found that he was losing power, so he came back to this country.

The lumbar region was extremely tender, but there was no deformity. Loss of motion and sensation were complete in the lower extremities, and of the bladder and rectum. Three vertebræ were diseased.

Mr D—, aged twenty-eight, was thrown from his horse in Natal on to his back. After some few minutes he was able to remount, and ride many miles to his home. His back was never free from pain, so he came over to this country. The fifth, sixth, and seventh dorsal vertebræ were diseased, but there was no deformity. Percussion induced great pain, but otherwise there was none. When sitting or standing there was no irregularity at the seat of disease, but increased convexity and extraordinary mobility.

Mr N—, aged twenty-six, was riding when his horse stumbled in going down a steep hill. He was much shaken,

and felt considerable pain in the neck. He continued his ride for several miles, and as he was making a tour, he continued it on foot. During the whole of this time he had pain in the neck. Eight months after the accident the fourth dorsal vertebra became prominent, and two months later motion of the lower limbs was impaired, and control over the bowel was lost. In the following month the intercostal muscles were paralysed. There was found, exactly one year after the accident, caries of the fourth and fifth dorsal vertebræ, and the intervertebral cartilage between them was destroyed. A portion of the body of the fourth dorsal vertebra was detached, and the medulla was compressed and softened to the extent of one inch.

When spinal disease is recognised in an early stage, whether in the child or in the adult, and treated as any other joint disease, and promptly, namely, with absolute rest, there is no reason why it should not terminate favorably; and there is every reason why it should not be succeeded by deformity, abscess, and paralysis. Disease may be cut short while in the inflammatory stage, and it is little liable to become chronic. And even when a vertebra first becomes slightly prominent, if absolute rest of the entire body is enforced, it is possible that caries may not extend further. When caries is established, especially in the dorsal region, some prominence of the spinous processes will remain.

There is no more doubt as to the result of such a case, when disease is recognised in an early stage and carefully treated, than after a similar injury to any other articulation. And it is perfectly known that the use of a joint may be entirely restored, even after serious injury, if complete immobility is for a time maintained, and the joint itself, as well as the limb, is adequately supported.

When the bodies of certain vertebræ have been destroyed, and a chasm has been produced, bony ankylosis may take place, even though three or four vertebræ are implicated in the disease; and our museums show that this may take place without deformity, and without

encroaching on the medullary canal. Specimens are not wanting which show this result of bony union even in the neck. In the lumbar region it is quite common, and in the dorsal region complete bony union takes place, but rarely without some deformity.

Paraplegia with loss of motion is due to pressure somewhat suddenly applied to the anterior column of the cord. And when sensation is also lost deeper seated injury has been incurred. This may be due to disorganisation, or it may be due to pressure which can be relieved, as in the following case, which I lately saw at Woodford :

A clergyman's daughter, aged fifteen, fell and struck her back in 1879. The accident was scarcely noticed, but some pain remained at the upper part of the back, and after seven months there was observed a prominence of some of the vertebræ. She was made to lie down, but was not otherwise treated. Suddenly she resolved to sit up, and forthwith, whilst she was exclaiming that she could sit up perfectly, was paralysed. I saw her on the following day, November, 1881. There was disease of the last cervical and two upper dorsal vertebræ, with some slight tenderness on moderate pressure, and some little deformity ; and there was total loss of motion of the lower extremities, and of sensation as high up as the ribs, and paralysis of the sphincters. She was kept perfectly motionless and flat on her back, and in four months, namely, in March, 1882, she had recovered both sensation and motion, and the control of the bladder and the rectum. During the period of paralysis she was greatly troubled by jerking of the limbs during sleep, and as she was going to sleep. This was so great that the legs were sometimes jerked out of bed.

CHAPTER VIII.

TREATMENT.

FROM what has gone before it is obvious that spinal disease may be divided into three stages; and for purposes of treatment it is convenient to consider:

1st. The acutely painful state.

2ndly. The painful state with rigidity.

3rdly. The state of deformity.

First. Immediately after the accident there is a period of acute pain, when it is well to apply an ice-bag to the injured part of the spine. And as the child throws himself about, being in acute pain, the ice should be pounded, otherwise it might injure him still further. At the same time a sufficient opiate may be given to induce sleep. As the pain becomes less acute cold applications may be discontinued, and an accurately fitting splint applied. This splint should be worn in the recumbent position until pain and tenderness have entirely ceased.

When a child continues to complain of pain after an accident, but is unable to define the seat of pain; when he no longer cares to play with other children as before, but lies down, and prefers to lie down, it should at once be suspected that he has met with an accident which needs attention. The injury may be so severe that it may lead to angular curvature of the spine; or the dorsal pain which has been excited may yield to a few weeks of recumbency and entire rest. Whether the accident is to terminate in angular curvature or whether the patient is to be restored after some few weeks to his former state of health and freedom from disease will depend entirely on the manner in which the accident is treated.

The patient should be placed perfectly flat on a well made horse-hair mattress, and without a pillow for the head; but the neck should be supported by a roll of linen immediately below the occiput, and a splint should be applied the whole length of the spine to give it adequate support. It is not important of what material the splint is constructed, so long as it fits accurately and the patient can lie in it comfortably.

The chief points to be desired are that it shall be made rapidly and easily; and, as regards the patient, that there shall be absence of pain, and a motionless condition.

When all pain has subsided, and the child can move freely on the bed without a splint, a starch bandage or a plaster of Paris bandage may be applied, and he may be allowed to move about the room. The bandage should be applied in the recumbent position. Suspension should never be permitted.

If disease were recognised and treated at this stage, we should never see angular curvature of the spine.

Secondly. When disease has progressed beyond the acute stage, and the patient is again able to move about, he controls the pain by steadying the body, either throwing the muscles of the back into action and thus making himself preternaturally erect, or by resting one or both hands on his thighs. Thus he moves about gently; no longer springing or jumping or stepping boldly, but he moves slowly along the floor; or, in coming down stairs, he plants both feet on every stair, with evident anxiety not to jar or shake the body. On examining the back, the muscles are found to be rigid to keep the spine fixed, and upright. And whatever he does, whether kneeling or sitting or standing, the muscles are thrown into action to prevent a sudden movement of the spine: he lowers himself on to his knee to pick up anything from the floor, and in lying down he is not less cautious. If the muscles are for a moment relaxed incautiously, the child screams with pain.

During all this time the back should be supported

by a splint accurately moulded to the surface, and the patient should be constantly recumbent. If the cervical vertebræ are injured the neck should be supported by the splint, and there should also be a prolongation from it to the occiput, to give support to the head and to prevent any movement upon the axis. Disease in this portion of the spine is much more serious than when it occurs lower down. Deformity is less than in the dorsal region, but softening is liable to take place in the upper part of the cord, and to terminate fatally. Also, the odontoid process may become displaced through destruction of the ligamentous attachments, and being displaced backwards, through a sudden movement of the head, will crush the medulla. Death, under these circumstances, is instantaneous.

Before an abscess has formed in the neck attention will have been attracted to the part by the pain which is suffered. And in the dorsal region there will probably be an angular projection before any abscess is discoverable. But when the lumbar vertebræ are diseased deformity shows itself later, and an abscess may first form and appear outside the erector spinæ.

But although abscess may have formed in connection with the upper dorsal vertebræ, showing itself perhaps in the back, piercing the muscles in close proximity to the spinous processes or along a rib; or though it be a lumbar abscess, or a psoas or an iliac abscess, it may be absorbed, and recovery may take place with a firmly ankylosed spine, and with little or no deformity. When, however, an abscess has formed in connection with the cervical vertebræ there is always a certain amount of shortening of the neck, even though there may be no angular projection. After an abscess has formed, unless a motionless condition of the patient is observed with absolute recumbency, it will not be absorbed.

When the abscess has to be evacuated the patient's life is in peril. It is therefore of prime importance to prevent the formation of abscess. The erect position

excites movements of the carious bones one upon another, and thus causes an increased formation of pus, which, indeed, may probably become so plentiful that it cannot be absorbed. Besides, it favours the gravitation of pus towards the groin. For these reasons, also, the recumbent position should be maintained.

Absolute rest in the recumbent position should be maintained until inflammation has ceased and repair is advanced.

Whilst a child is recumbent the general health improves: he sleeps and eats well, and is happy, all his functions are well performed, there is no fear of abscess, and repair is progressing. But when he is allowed to get up he suffers with pain of various kinds, "rheumatic" and "growing pains;" he becomes restless in sleep, loses appetite, colour, and weight, notwithstanding the bandage or support that is applied.

Until ankylosis is advanced movement should not be permitted.

When a child has been uncared for, and has been allowed to move about, supporting himself by resting his hands on the thighs, any form of splint or bandage around the body affords immense relief; and thus bound up of course he will walk upright, and without resting his hands on his thighs. It has always been a difficulty in hospital practice after the support has been effectively applied, and the child is relieved from pain, to keep him recumbent. He has no difficulty in sitting or standing, and he does it when the nurse is absent. But the question which every surgeon has to put to himself is, Is it right to allow him to sit or stand?

An excellent surgeon, than whom there was none better, Prof. Pirrie, answered this question thus:—"Rest of the diseased parts and the recumbent position, whether the body be prone or supine, are of the utmost importance from the very commencement of the disease, until a cure is effected by ankylosis."*

* 'Principles of Surgery,' 2nd edition, p. 463.

But it is not sufficient that the child shall be kept recumbent. It is of the utmost importance that he should be kept recumbent until repair takes place, so that abscess and deformity may not occur; but unless a retentive apparatus be added to the recumbent position abscess and deformity will surely follow. It matters very little what the form of support is so long as it is comfortable and holds the spine perfectly in position. A support to the back is even more necessary than in fracture of a long bone.

Growth is sometimes rapid, after having been checked, when recumbency is first observed. It is not unusual to find that disease having gone on for one, two, or three years growth has stopped, almost, if not entirely, during this period; but with recumbency it recommences, and the lost height is regained in the course, perhaps, of two years. Thus, I have known three inches in height to be gained in six months in a case where during the previous three years there had been absolutely no growth, and this continued until the full height was made up.

When the recumbent position is maintained there is small difficulty with appetite, and the patient is ready to receive whatever is given. The diet should consist mainly of animal substances, and there is no necessity to feed inordinately. The tendency is under these circumstances to increase in weight, and sometimes this takes place to a very inconvenient extent, so that it becomes necessary to observe a strict regimen. This is especially necessary with adults.

Thirdly. When deformity arises is it to be removed? Is it justifiable to remove the angular deformity, and straighten the spine?

It has been so long taught that the correct treatment of spinal disease is to allow the disjoined vertebræ to fall one upon another, and thus to become in some sort united, that many hold the opinion that an angular curvature cannot be removed. Mr Solly even wrote that only an ignorant man or a knave would pretend to correct the deformity of angular curvature, and he contends

“that the maintenance of the curve, and not its obliteration, is the thing to be sought for.”* But Mr Solly’s views of treatment are also peculiar. He says, “In all cases of angular curvature entire rest is absolutely necessary, and this must be continued for some time, seldom less than four or five months, and often for a year or a year and a half. The exact position in bed must vary with circumstances: as a general rule that which is easiest to the patient is best, and the lateral position will frequently be found so, taking care to support the back with pillows that are fixed so that they cannot be pushed away.” And Prof. Bauer says “we are able to arrest the progress of the disease, and we can avert, but cannot remove its ulterior consequences.”†

The removal of angular deformity depends on the stage at which disease has arrived. Until repair has commenced, until ankylosis, true or spurious, has formed, angular deformity is removable.

Before there is any attempt at solidification, angular deformity may be removed; but when repair has commenced it is not justifiable to interfere with it.

For the purposes of treatment the prone position is perhaps the best; but it is advisable that the patient shall be placed supine for several hours in the twenty-four. Any change that is effected in the spine can only be done very gradually, and it is made simply by observing the horizontal position and by the use of splints.

In the first instance a cast of the back should be taken, and upon this a splint may be blocked to fit accurately to the back, and to support efficiently the ribs. In this the patient will lie. It will be removed for purposes of cleanliness when needed. After a time it will be observed that the splint no longer fits well, and that it needs to be reblocked; and perhaps this may have to be done every three or four months or even more frequently, until there is little or no projection left. Abscess becomes absorbed

* ‘Surgical Experiences,’ pp. 27 and 77.

† *Op. cit.*, p. 134.

during this period, and ankylosis eventually takes place. When bony union has formed, a spinal support to give additional security may be applied, and exercise may be permitted.

A good form of support is made with a horse-shoe shaped pad. This will support the ribs, without making pressure on the spinous processes. It should be so placed that if there be an angular projection remaining the pad shall be immediately below it, and it may rest on the angles of the ribs. When deformity is low down the pad may be borne on a single lever; but in the upper dorsal region a double lever will be necessary. And when the neck is the seat of disease, the support should extend to the occiput, so that no motion whatever of the head or of the cervical vertebræ may take place.

Dr Taylor's antero-posterior support is useful when the figure is slight, and equally so is Mr Chance's modification of it. The first-named support is more solid, and better adapted for wear, and I therefore generally employ it.

Exercise may be permitted when repair has commenced, if a well fitting apparatus is worn; but violent exercise should not be permitted until consolidation is complete.

It is difficult to persuade patients and parents that care is needed for many months after repair is far advanced.

A child was sent over from China with disease of the ninth dorsal vertebra. He was a puny delicate boy, and had been half starved on the voyage, and the chances were greatly against life. However, he was kept recumbent in a splint, and eventually grew into a fine, strong and tall lad, and at ten years of age, he could bend about in every possible manner. He went to school at Harrow, and walked a mile to and fro daily. He was in wonderful health and spirits, and ready for any fun. He was cautioned that all his movements should be even and regular, and that no wrestling or tumbling or rough play of any kind could be allowed. For fully one year this

kind of life had been led, when on an unlucky day he spied a wood-pigeon's nest in a high tree in his father's grounds. The next moment he was on his way up to it, but he did not arrive there ; for just before reaching it he nearly fell. He broke through some bony union, and this was followed by inflammation and many abscesses, which at length terminated fatally.

It is important to support and protect the spine, and especially the seat of disease, long after a patient has come to enjoy freedom of movement.

Spinal abscess, in whatever position, should not be punctured, unless tension is so great, that ulceration is imminent ; for it is better that the pus shall be absorbed than that the abscess shall be opened. When the abscess has been opened and is discharging, the pus is apt to assume an unhealthy character, and even to become foetid. There is then danger of septicæmia, even though the opening may have been made antiseptically : hectic may then supervene.

Under these circumstances, the abscess should be freely laid open, and the cavity washed out with a weak carbolic acid lotion.

Although an abscess should never be opened unnecessarily, an opening should never be allowed to form, through ulceration. First, the most convenient spot will be selected for the opening ; but this is very frequently not selected by nature. Even it might be said that the most convenient situation is very seldom selected by nature ; and a second and more dependent opening has to be made. A portion of the contents of the abscess can be drawn off with the aspirator, tension may be relieved, and the wound can be closed, whereas, if ulceration takes place, the wound remains open. This withdrawal of pus may be repeated if necessary. But if the discharge is thick and flaky, and will not pass through the needle of the aspirator, the opening may be enlarged antiseptically and the contents of the abscess withdrawn. The aspirator needles should be made of fine steel and

with finer points and much larger eyes, and the needle itself of larger calibre than is generally in use. It is a perfect mode of opening an abscess when the puncture can be made readily and the contents of the abscess drawn off in a stream. In two days the little wound has entirely closed ; and there is no fear with this instrument, of incurring the danger of foetid pus.

In spinal abscess an open wound is to be avoided and drainage tubes are to be shunned. Few things are more injurious and debilitating than a long continued, exhausting discharge. If it were only for cleanliness the aspirator is to be recommended. But this is the least of its virtues.

When pus having collected in considerable quantity, is found to be burrowing, it should be withdrawn without delay.

I lately saw a young Jew, who had caries of the fourth and fifth dorsal vertebræ, and an iliac abscess. He was moving about without support and resting his hand on the thigh. Pus having burrowed, had passed through the inguinal canal into the scrotum, and also the perinæum was infiltrated with pus ; so that both the perinæum and the scrotum had to be opened.

In conclusion I may venture thus to sum up, that so long as inflammation exists recumbency and support to the diseased spinal column are to be observed ; and when angular projection with or without abscess has taken place still recumbency in the prone and supine positions alternately, together with a support is to be observed ; care being taken to alter and lengthen the support as growth takes place. And when ankylosis has taken place exercise is to be permitted with adequate support to the back.

Now, that recumbency has been so much insisted on it will perhaps be asked, when then is suspension to be employed ?

Suspension would scarcely be employed, unless there were deformity, for the object of suspension is to remove or diminish angular curvature. Sometimes Dr Sayro

practises extension thus : a boy, five years of age, with Pott's disease in the lumbar region "was held out as straight as possible by two assistants pulling, one from the shoulders, and the other from the ankles," and the straightening of the spine thus produced, is shown in Fig. 5*. But Dr Sayre prefers suspension to stretching the spine, and he writes thus : "As it is difficult for an assistant to hold a patient suspended during the application of the dressing, I make use of a very convenient apparatus contrived by Mr Reynders, of New York, which consists of a curved iron cross-beam, to which is attached an adjustable head and chin collar with straps fitted to axillary bands. To a hook in the centre is fixed a compound pulley, the other end of which is secured either to a hook in the ceiling or to the top of an iron tripod about ten feet in height. The head and chin collar and the axillary supports having been carefully adjusted, the patient is gradually drawn up until the feet swing just clear of the floor."† The plaster-jacket is then applied, and Dr Sayre finds the following result ; that, "as soon as the plaster-jacket has become hardened, the patient has been able to walk about and *in many instances, where paralysis had been present before the jacket was applied, the sufferer had been able to walk without difficulty after the application was made.*"*

This is perhaps the most extraordinary statement that I ever read. With the converse of this most of us are familiar, viz. that a patient having the full use of his limbs may be suspended and let down utterly paralysed. Unfortunately not in London only but elsewhere this result is not uncommon. But it is a miracle and otherwise unknown to science, that paraplegia may be removed by suspension in the course of half an hour.

The pathological conditions in which suspension is desirable in spinal disease have not been laid down. Nor do I find in Dr Sayre's work any mention of the pathological conditions which are considered favorable or un-

* 'Spinal Disease and Spinal Curvature,' p. 25.

† Op. cit., pp. 17 and 19.

favorable for his treatment by suspension. I have introduced three woodcuts which were done by Mr Bagg years ago, and which I now use for the first time (Figs. 31, 32, and 33). They represent appearances which are familiar to every pathologist.

It would never occur to me to suspend cases where there was such disease present as is there represented ; but when it is not defined what class of cases should be suspended, every case, of whatever age, from some few months to seventy years, with acute disease and with ankylosis, spurious and true, is in danger of being suspended. And it is not alone that the danger exists for such patients, but suspension is practised at every age and in every stage of disease, and the result is that more severe cases are now seen than I ever remember to have seen before suspension was in vogue.

Suspension was practised by Glisson in the seventeenth century. When, however, pathology came to be better taught such treatment was held to be barbarous. Except that such treatment is, according to Dr Sayre, an admirable mode of removing paralysis, one would really suppose that no system of treatment could be more unjustifiable. But we are told that in parts of Germany they are not content simply with suspending their patients by the head and shoulders, but in this position they administer chloroform, so that, by removing muscular action, the straightening of the spine may be more effective. I have not yet heard of this practice being carried out in England.

For my own part I should not, under any circumstances, suspend a patient with caries of the spine, not even to remove paralysis ; first, because it is attended with extreme danger, and secondly, because it is entirely unnecessary. It is dangerous, inasmuch as it induces abscess and paralysis, and it breaks through repair ; and it is unnecessary because the horizontal position effects everything that can be desired without danger. The horizontal position promotes union also, whereas suspension prevents it.

With regard to the second part of the process, the

objection that I take to the use of the swathing bands impregnated with plaster of Paris is that the jacket harbours vermin to an extraordinary extent, and further that the same effect may be produced in a more simple and a more cleanly manner, viz. with a movable splint. And whether this splint be made of felt, leather, or poroplastic material, it should be made to fit exactly to the back in every part, and thus fitting it matters very little of what the splint is constructed. Perhaps the poroplastic splint is easier to make than any other, but it seldom fits well. But the most serious condemnation of the jacket is offered by Dr Sayre himself, in the following words :

“ When the thorax is thus firmly secured,” he writes, “ the anus and the perinæum will rise and fall synchronously with the diaphragm, and the respiration be carried on without difficulty, so long as these parts are free from pressure. If manual pressure be applied to the perinæum a feeling of suffocation is produced. *It is therefore necessary in some cases, when the thorax is thus secured, that the patient should sit upon a chair with a hole in the seat, like a close-stool.*”*

Thus the jacket prevents thoracic respiration and promotes collapse of the walls of the chest; and its effect on a deformed thorax is more injurious than the tightest stays.

Again, Sayre’s “ jury-mast ” is the head-rest of Shel-drake : a frightful piece of machinery, which years ago I have seen in use. For twenty years it has not been in use, because a less unsightly, but equally effective, apparatus is now worn.

My colleague, Mr Baker, has put the case of suspension and the employment of the jacket well. He says, “ In the consideration of this subject in relation to lateral curvature of the spine it appears strange that the same form of treatment that is advised for the cure of caries of the vertebræ should have been so confidently recommended for the treatment of lateral curvature, as the two affections are so

* Op. cit., p. 12.

entirely different in their pathology and progress. Angular curvature is always the result of one cause, viz. caries of the vertebræ, so that it is rational to treat all cases in a similar manner; on the other hand, lateral curvature is due to a variety of causes, and it appears somewhat irrational to suggest one form of treatment for all cases, and for all the different stages through which the deformity passes." *

Suspension in caries of the spine, whether in the inflammatory stage or when repair is taking place, is opposed to every notion of pathology, and should be prohibited. And although the plaster jacket may give relief where no previous application has been made, and is therefore fitted for outcasts or those living beyond other help, it is unfit to be used in a civilised society, and where we pride ourselves on caring for our poor. No cure is possible with such a bandage without recumbency, and with recumbency the bandage is not desirable. It has been employed without consideration, and the sooner its evils are understood the better it will be for patients who need some form of support.

* "On the treatment of Spinal Curvature, with special reference to Sayre's method." 'Transactions of the International Medical Congress,' 1881, vol. iv, p. 165.

Catalogue B]

London, 11, New Burlington Street
March, 1883.

SELECTION

FROM

J. & A. CHURCHILL'S GENERAL CATALOGUE

COMPRISING

ALL RECENT WORKS PUBLISHED BY THEM

ON THE

ART AND SCIENCE OF MEDICINE



N.B.—As far as possible, this List is arranged in the order in which medical study is usually pursued.

J. & A. CHURCHILL publish for the following Institutions
and Public Bodies :—

H.M. STATIONERY OFFICE.

VIVISECTION FORMS AND CERTIFICATES.

A to F (6 at $\frac{1}{2}$ d. each). Application for Licence, $\frac{1}{2}$ d.

ROYAL COLLEGE OF SURGEONS.

CATALOGUES OF THE MUSEUM.

Nineteen separate Catalogues (List and Prices can be obtained of J. & A. CHURCHILL).

GUY'S HOSPITAL.

REPORTS BY THE MEDICAL AND SURGICAL STAFF.

The last Volume (XXV., Third Series) was issued in 1881. Price 7s. 6d.

LONDON HOSPITAL.

PHARMACOPŒIA OF THE HOSPITAL. 3s.

CLINICAL LECTURES AND REPORTS BY THE MEDICAL AND
SURGICAL STAFF. Vols. I. to IV., 7s. 6d. each.

ST. BARTHOLOMEW'S HOSPITAL.

CATALOGUE OF THE ANATOMICAL AND PATHOLOGICAL
MUSEUM. Vol. I.—Pathology. 15s.

ST. GEORGE'S HOSPITAL.

REPORTS BY THE MEDICAL AND SURGICAL STAFF.

The last Volume (X.) was issued in 1880. Price 7s. 6d.

CATALOGUE OF THE PATHOLOGICAL MUSEUM. 15s.

SUPPLEMENTARY CATALOGUE (1882). 5s.

ST. THOMAS'S HOSPITAL.

REPORTS BY THE MEDICAL AND SURGICAL STAFF.

Annually. Vol. XI., New Series (1882). 7s. 6d.

ROYAL LONDON OPHTHALMIC HOSPITAL.

REPORTS BY THE MEDICAL AND SURGICAL STAFF.

Occasionally. Vol. X., Part III. (August, 1882). 5s.

MEDICO-PSYCHOLOGICAL ASSOCIATION.

JOURNAL OF MENTAL SCIENCE.

Quarterly. Price 3s. 6d. each, or 14s. per annum.

PHARMACEUTICAL SOCIETY OF GREAT BRITAIN.

PHARMACEUTICAL JOURNAL AND TRANSACTIONS.

Each Week. Price 4d. each, or 20s. per annum, post free.

BRITISH PHARMACEUTICAL CONFERENCE.

YEAR BOOK OF PHARMACY.

In December. Price 10s.

BRITISH DENTAL ASSOCIATION.

JOURNAL OF THE ASSOCIATION AND MONTHLY REVIEW
OF DENTAL SURGERY.

On the 15th of each Month. Price 6d., or 13s. per annum, post free.

A SELECTION

FROM

J. & A. CHURCHILL'S GENERAL CATALOGUE,

COMPRISING

ALL RECENT WORKS PUBLISHED BY THEM ON THE
ART AND SCIENCE OF MEDICINE.

N.B.—*J. & A. Churchill's Descriptive List of Works on Chemistry, Materia Medica, Pharmacy, Botany, Photography, Zoology, the Microscope, and other Branches of Science, can be had on application.*

Practical Anatomy:

A Manual of Dissections. By CHRISTOPHER HEATH, Surgeon to University College Hospital. Fifth Edition. Crown 8vo, with 24 Coloured Plates and 269 Engravings, 15s.

Wilson's Anatomist's Vade-

Mecum. Tenth Edition. By GEORGE BUCHANAN, Professor of Clinical Surgery in the University of Glasgow; and HENRY E. CLARK, M.R.C.S., Lecturer on Anatomy at the Glasgow Royal Infirmary School of Medicine. Crown 8vo, with 450 Engravings (including 26 Coloured Plates), 18s.

Braune's Atlas of Topographi-

cal Anatomy, after Plane Sections of Frozen Bodies. Translated by EDWARD BELLAMY, Surgeon to, and Lecturer on Anatomy, &c., at, Charing Cross Hospital. Large Imp. 8vo, with 34 Photolithographic Plates and 46 Woodcuts, 40s.

An Atlas of Human Anatomy.

By RICKMAN J. GODLEE, M.S., F.R.C.S., Assistant Surgeon and Senior Demonstrator of Anatomy, University College Hospital. With 48 Imp. 4to Plates (112 figures), and a volume of Explanatory Text, 8vo, £4 14s. 6d.

Surgical Anatomy:

A Series of Dissections, illustrating the Principal Regions of the Human Body. By JOSEPH MACLISE. Second Edition. 52 folio Plates and Text. Cloth, £3 12s.

Medical Anatomy.

By FRANCIS SIBSON, M.D., F.R.C.P., F.R.S. Imp. folio, with 21 Coloured Plates, cloth, 42s., half-morocco, 50s.

Anatomy of the Joints of Man.

By HENRY MORRIS, Surgeon to, and Lecturer on Anatomy and Practical Surgery at, the Middlesex Hospital. 8vo, with 44 Lithographic Plates (several being coloured) and 13 Wood Engravings, 16s.

Manual of the Dissection of the Human Body.

By LUTHER HOLDEN, Consulting Surgeon to St. Bartholomew's and the Foundling Hospitals, and JOHN LANGTON, F.R.C.S., Surgeon and Lecturer on Anatomy at St. Bartholomew's Hospital. Fourth Edition. 8vo, with 170 Engravings, 16s.

By the same Author.

Human Osteology:

Sixth Edition, edited by the Author and JAMES SHUTER, F.R.C.S., M.A., M.B., Assistant Surgeon to St. Bartholomew's Hospital. 8vo, with 61 Lithographic Plates and 89 Engravings. 16s.

Also.

Landmarks, Medical and Surgical.

Third Edition. 8vo, 3s. 6d.

The Student's Guide to Surgical

Anatomy: An Introduction to Operative Surgery. By EDWARD BELLAMY, F.R.C.S. and Member of the Board of Examiners. Fcap. 8vo, with 76 Engravings, 7s.

The Student's Guide to Human

Osteology. By WILLIAM WARWICK WAGSTAFFE, late Assistant Surgeon to St. Thomas's Hospital. Fcap. 8vo, with 23 Plates and 66 Engravings, 10s. 6d.

The Anatomical Remembrancer; or, Complete Pocket Anatomist. Eighth Edition. 32mo, 3s. 6d.

Diagrams of the Nerves of the Human Body, exhibiting their Origin, Divisions, and Connections, with their Distribution to the Various Regions of the Cutaneous Surface, and to all the Muscles. By WILLIAM H. FLOWER, F.R.C.S., F.R.S., Hunterian Professor of Comparative Anatomy to the Royal College of Surgeons. Third Edition, with 6 Plates. Royal 4to, 12s.

Atlas of Pathological Anatomy. By Dr. LANCEREAUX. Translated by W. S. GREENFIELD, M.D., Professor of Pathology in the University of Edinburgh. Imp. 8vo, with 70 Coloured Plates, £5 5s.

A Manual of Pathological Anatomy. By C. HANDFIELD JONES, M.B., F.R.S.; and EDWARD H. SIEVEKING, M.D., F.R.C.P. Edited (with considerable enlargement) by J. F. PAYNE, M.D., F.R.C.P., Lecturer on General Pathology at St. Thomas's Hospital. Second Edition. Crown 8vo, with 195 Engravings, 16s.

Lectures on Pathological Anatomy. By SAMUEL WILKS, M.D., F.R.S.; and WALTER MOXON, M.D., Physician to Guy's Hospital. Second Edition. 8vo, Plates, 18s.

Post-Mortem Examinations: A Description and Explanation of the Method of performing them, with especial reference to Medico-Legal Practice. By Prof. VIRCHOW. Translated by Dr. T. P. SMITH. Second Edition. Fcap. 8vo, with 4 Plates, 3s. 6d.

The Human Brain: Histological and Coarse Methods of Research. A Manual for Students and Asylum Medical Officers. By W. BEVAN LEWIS, L.R.C.P. Lond., Deputy Medical Superintendent to the West Riding Lunatic Asylum. 8vo, with Wood Engravings and Photographs, 8s.

Principles of Human Physiology. By W. B. CARPENTER, C.B., M.D., F.R.S. Ninth Edition. By HENRY POWER, M.B., F.R.C.S. 8vo, with 3 Steel Plates and 377 Wood Engravings, 31s. 6d.

A Treatise on Human Physiology. By JOHN C. DALTON, M.D., Professor in the College of Physicians and Surgeons, New York. Seventh Edition. Med. 8vo, with 252 Engravings, 20s.

Text-Book of Physiology. By J. FULTON, M.D., Professor of Physiology, &c., in Trinity Medical College, Toronto. Second Edition. 8vo, with 152 Engravings, 15s.

Sanderson's Handbook for the Physiological Laboratory. By E. KLEIN, M.D., F.R.S.; J. BURDON-SANDERSON, M.D., F.R.S.; MICHAEL FOSTER, M.D., F.R.S.; and T. LAUDER BRUNTON, M.D., F.R.S. 8vo, with 123 Plates, 24s.

Histology and Histo-Chemistry of Man. By HEINRICH FREY, Professor of Medicine in Zurich. Translated by ARTHUR E. J. BARKER, Assistant-Surgeon to University College Hospital. 8vo, with 608 Engravings, 21s.

The Marriage of Near Kin, Considered with respect to the Laws of Nations, Results of Experience, and the Teachings of Biology. By ALFRED H. HUTH. 8vo, 14s.

Medical Jurisprudence: Its Principles and Practice. By ALFRED S. TAYLOR, M.D., F.R.C.P., F.R.S. Third Edition, by THOMAS STEVENSON, M.D., F.R.C.P., Lecturer on Medical Jurisprudence at Guy's Hospital. 2 vols. 8vo, with Engravings.

By the same Author.

A Manual of Medical Jurisprudence. Tenth Edition. Crown 8vo, with 55 Engravings, 14s.

Also.

Poisons, In Relation to Medical Jurisprudence and Medicine. Third Edition. Crown 8vo, with 104 Engravings, 16s.

Lectures on Medical Jurisprudence. By FRANCIS OGSTON, M.D., Professor in the University of Aberdeen. Edited by FRANCIS OGSTON, JUN., M.D. 8vo, with 12 Copper Plates, 18s.

A Handy-Book of Forensic Medicine and Toxicology. By C. MEYMOTT TIDY, M.D., F.C.S., and W. BATHURST WOODMAN, M.D., F.R.C.P. 8vo, with 8 Lithographic Plates and 116 Engravings, 31s. 6d.

Microscopical Examination of Drinking Water. By JOHN D. MACDONALD, M.D., F.R.S., Assistant Professor in the Army Medical School. 8vo, with 24 Plates, 7s. 6d.

Sanitary Examinations Of Water, Air, and Food. A Vade-Mecum for the Medical Officer of Health. By CORNELIUS B. FOX, M.D., F.R.C.P. Crown 8vo, with 94 Engravings, 12s. 6d.

Sanitary Assurance: A Lecture at the London Institution. By Prof. F. DE CHAUMONT, F.R.S. With Short Addresses by J. E. ERICHSEN, F.R.S., Sir J. FAYRER, K.C.S.I., and R. BRUDENELL CARTER, F.R.C.S., &c. Royal 8vo, 1s.

A Manual of Practical Hygiene.

By E. A. PARKES, M.D., F.R.S. Sixth Edition. By F. DE CHAUMONT, M.D., F.R.S., Professor of Military Hygiene in the Army Medical School. 8vo, with Plates and Engravings. (*In the Press.*)

Dangers to Health:

A Pictorial Guide to Domestic Sanitary Defects. By T. PRIDGIN TEALE, M.A., Surgeon to the Leeds General Infirmary. Third Edition. 8vo, with 70 Lithograph Plates (mostly coloured). 10s.

A Handbook of Hygiene and Sanitary Science.

By GEO. WILSON, M.A., M.D., Medical Officer of Health for Mid-Warwickshire. Fourth Edition. Post 8vo, with Engravings, 10s. 6d.

Also.

Healthy Life and Healthy Dwellings:

A Guide to Personal and Domestic Hygiene. Fcap. 8vo, 5s.

Contributions to Military and State Medicine.

By JOHN MARTIN, L.R.C.S.E., Surgeon Army Medical Department. 8vo, 10s. 6d.

Pay Hospitals and Paying Wards throughout the World.

By HENRY C. BURDETT, late Secretary to the Seamen's Hospital Society. 8vo, 7s.

By the same Author.

Cottage Hospitals — General, Fever, and Convalescent:

Their Progress, Management, and Work. Second Edition, with many Plans and Illustrations. Crown 8vo, 14s.

Dress: Its Sanitary Aspect.

A Paper read before the Brighton Social Union, Jan. 30, 1880. By BERNARD ROTH, F.R.C.S. 8vo, with 8 Plates, 2s.

Manual of Anthropometry:

A Guide to the Measurement of the Human Body, containing an Anthropometrical Chart and Register, a Systematic Table of Measurements, &c. By CHARLES ROBERTS, F.R.C.S. 8vo, with numerous Illustrations and Tables, 6s. 6d.

Madness:

In its Medical, Legal, and Social Aspects. Lectures by EDGAR SHEPPARD, M.D., M.R.C.P., Professor of Psychological Medicine in King's College. 8vo, 6s. 6d.

Idiocy and Imbecility.

By WILLIAM W. IRELAND, M.D., Medical Superintendent of the Scottish National Institution for the Education of Imbecile Children at Larbert, Stirlingshire. 8vo, with Engravings, 14s.

A Manual of Psychological

Medicine: With an Appendix of Cases. By JOHN C. BUCKNILL, M.D., F.R.S., and D. HACK TUKE, M.D., F.R.C.P. Fourth Edition. 8vo, with 12 Plates (30 Figures) and Engravings, 25s.

The Student's Guide to the Practice of Midwifery.

By D. LLOYD ROBERTS, M.D., F.R.C.P., Physician to St. Mary's Hospital, Manchester. Second Edition. Fcap. 8vo, with 111 Engravings, 7s.

Handbook of Midwifery for Midwives:

from the Official Handbook for Prussian Midwives. By J. E. BURTON, L.R.C.P. Lond., Senior Assistant Medical Officer, Ladies' Charity, &c., Liverpool. With Engravings. Fcap. 8vo, 6s.

Lectures on Obstetric Operations:

Including the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labour. By ROBERT BARNES, M.D., F.R.C.P., Obstetric Physician to St. George's Hospital. Third Edition. 8vo, with 124 Engravings, 18s.

By the same Author.

A Clinical History of Medical and Surgical Diseases of Women.

Second Edition. 8vo, with 181 Engravings, 28s.

West on the Diseases of Women.

Fourth Edition, revised by the Author, with numerous Additions by J. MATTHEWS DUNCAN, M.D., F.R.C.P., Obstetric Physician to St. Bartholomew's Hospital. 8vo, 16s.

Observations on the Cæsarean Section, Craniotomy, and on other Obstetric Operations, with Cases.

By THOMAS RADFORD, M.D., late Consulting Physician, St. Mary's Hospital, Manchester. Second Edition, with Plates. 8vo, 10s.

Clinical Lectures on Diseases of Women:

Delivered in St. Bartholomew's Hospital, by J. MATTHEWS DUNCAN, M.D., F.R.C.P. Second Edition. 8vo, 14s.

By the same Author.

Papers on the Female Perineum, &c.

8vo, 6s.

The Principles and Practice of Gynæcology.

By THOMAS ADDIS EMMET, M.D., Surgeon to the Woman's Hospital, New York. Second Edition. Royal 8vo, with 133 Engravings, 24s.

Diseases of the Uterus, Ovaries, and Fallopian Tubes:

A Practical Treatise by A. COURTY, Professor of Clinical Surgery, Montpellier. Translated from Third Edition by his Pupil, AGNES McLAREN, M.D., M.K.Q.C.P.I., with Preface by J. MATTHEWS DUNCAN, M.D., F.R.C.P. 8vo, with 424 Engravings, 24s.

- The Student's Guide to the Diseases of Women.** By ALFRED L. GALABIN, M.D., F.R.C.P., Assistant Obstetric Physician to Guy's Hospital. Second Edition. Fcap. 8vo, with 70 Engravings, 7s. 6d.
- Notes on the Diseases of Women.** Specially designed for Students preparing for Examination. By J. J. REYNOLDS, M.R.C.S. Fcap. 8vo, 2s. 6d.
By the same Author.
- Notes on Midwifery:**
Specially designed for Students preparing for Examination. Fcap. 8vo, 4s.
- Practical Gynæcology:**
A Handbook of the Diseases of Women. By HEYWOOD SMITH, M.D. Oxon., Physician to the Hospital for Women, &c. Second Edition. Crown 8vo, with Engravings. (*In the Press.*)
By the same Author.
- Dysmenorrhea, its Pathology and Treatment.** Crown 8vo, with Engravings, 4s. 6d.
- Obstetric Aphorisms:**
For the Use of Students commencing Midwifery Practice. By JOSEPH G. SWAYNE, M.D. Seventh Edition. Fcap. 8vo, with Engravings, 3s. 6d.
- Obstetric Medicine and Surgery:**
Their Principles and Practice. By F. H. RAMSBOTHAM, M.D., F.R.C.P. Fifth Edition. 8vo, with 120 Plates, 22s.
- A Complete Handbook of Obstetric Surgery.** Giving Short Rules of Practice in every Emergency. By CHARLES CLAY, late Surgeon to St. Mary's Hospital, Manchester. Third Edition. Fcap. 8vo, with 91 Engravings, 6s. 6d.
- Schroeder's Manual of Midwifery,** including the Pathology of Pregnancy and the Puerperal State. Translated by CHARLES H. CARTER, B.A., M.D. 8vo, with Engravings, 12s. 6d.
- Influence of Posture on Women** in Gynecic and Obstetric Practice. By J. H. AVELING, M.D., Physician to the Chelsea Hospital for Women. 8vo, 6s.
By the same Author.
- The Chamberlens and the Midwifery Forceps:** Memorials of the Family, and an Essay on the Invention of the Instrument. 8vo, with Engravings, 7s. 6d.
- A Handbook of Uterine Therapeutics,** and of Diseases of Women. By E. J. TILT, M.D., M.R.C.P. Fourth Edition. Post 8vo, 10s.
By the same Author.
- The Change of Life**
In Health and Disease: a Clinical Treatise on the Diseases of the Nervous System incidental to Women at the Decline of Life. Fourth Edition. 8vo, 10s. 6d.
- Ovarian and Uterine Tumours:**
Their Pathology and Surgical Treatment. By T. SPENCER WELLS, F.R.C.S., Consulting Surgeon to the Samaritan Hospital. 8vo, with Engravings, 21s.
- Rupture of the Female Perineum:** Its Treatment, immediate and remote. By GEORGE G. BANTOCK, M.D., Surgeon to the Samaritan Hospital. 8vo, with 2 Plates, 3s. 6d.
- Chronic Disease of the Heart:**
Its Bearings upon Pregnancy, Parturition, and Childbed. By ANGUS MACDONALD, M.D., F.R.S.E., Physician to the Edinburgh Royal Infirmary. 8vo, with Engravings, 8s. 6d.
- The Female Pelvic Organs,**
Their Surgery, Surgical Pathology, and Surgical Anatomy, in a Series of Coloured Plates taken from Nature: with Commentaries, Notes, and Cases. By HENRY SAVAGE, M.D., F.R.C.S., Consulting Officer of the Samaritan Free Hospital. Fifth Edition. Roy. 4to, with 17 Lithographic Plates (15 coloured) and 52 Woodcuts, £1 15s.
- Lectures on Diseases of the Nervous System,** especially in Women. By S. WEIR MITCHELL, M.D., Physician to the Philadelphia Infirmary for Diseases of the Nervous System. With 5 Plates. Post 8vo, 8s.
- A Treatise on the Diseases of Children.** For Practitioners and Students. By W. H. DAY, M.D., Physician to the Samaritan Hospital for Women and Children. Crown 8vo, 12s. 6d.
- The Wasting Diseases of Children.** By EUSTACE SMITH, M.D., Physician to the King of the Belgians, Physician to the East London Hospital for Children. Third Edition. Post 8vo, 8s. 6d.
By the same Author.
- Clinical Studies of Disease in Children.** Second Edition. Post 8vo. (*In the Press.*)
- Infant Feeding and its Influence on Life;** or, the Causes and Prevention of Infant Mortality. By C. H. F. ROUTH, M.D., Senior Physician to the Samaritan Hospital. Third Edition. Fcap. 8vo, 7s. 6d.
- A Practical Manual of the Diseases of Children.** With a Formulary. By EDWARD ELLIS, M.D. Fourth Edition. Crown 8vo, 10s.
By the same Author.
- A Manual of what every Mother should know.** Fcap. 8vo, 1s. 6d.
- Handbook for Nurses for the Sick.** By ZEPHERINA P. VEITCH. Second Edition. Crown 8vo, 3s. 6d.

A Manual of Nursing, Medical and Surgical. By CHARLES J. CULLINGWORTH, M.D., Physician to St. Mary's Hospital, Manchester. Fcap. 8vo, 3s. 6d.

A Manual for Hospital Nurses And others engaged in Attending on the Sick. By EDWARD J. DOMVILLE, Surgeon to the Exeter Lying-in Charity. Fourth Edition. Crown 8vo, 2s. 6d.

Notes on Fever Nursing. By J. W. ALLAN, M.B., Superintendent and Physician, Glasgow Fever Hospital. Crown 8vo, with Engravings, 2s. 6d.

Manual of Botany : Including the Structure, Functions, Classification, Properties, and Uses of Plants. By ROBERT BENTLEY, Professor of Botany in King's College and to the Pharmaceutical Society. Fourth Edition. Crown 8vo, with 1,185 Engravings, 15s.

Medicinal Plants : Being descriptions, with original figures, of the Principal Plants employed in Medicine, and an account of their Properties and Uses. By Prof. BENTLEY and Dr. H. TRIMEN. In 4 vols., large 8vo, with 306 Coloured Plates, bound in Half Morocco, Gilt Edges. £11 11s.

Royle's Manual of Materia Medica and Therapeutics. Sixth Edition, by JOHN HARLEY, M.D., Physician to St. Thomas's Hospital. Crown 8vo, with 139 Engravings, 15s.

A Manual of Practical Therapeutics. By E. J. WARING, C.I.E., M.D., F.R.C.P. Lond. Third Edition. Fcap. 8vo, 12s. 6d.

The Student's Guide to Materia Medica and Therapeutics. By JOHN C. THOROWGOOD, M.D., F.R.C.P. Second Edition. Fcap. 8vo, 7s.

Materia Medica and Therapeutics. By CHARLES D. F. PHILLIPS, M.D., late Lecturer on Materia Medica and Therapeutics at the Westminster Hospital Medical School.

Vol. 1—Vegetable Kingdom. 8vo, 15s.

Vol. 2—Inorganic Substances. 8vo, 21s.

Therapeutical Remembrancer. By JOHN MAYNE, M.D. Second Edition. 16mo, 3s. 6d.

By the same Author.

Notes on Poisons. Mounted and Varnished for the Surgery. 18 in. by 12 in. 1s. 6d.

Indian Notes. The Voyage out; Travelling in India; Upper India; Stations; The Hills; Mineral Waters; Herbs and Simples. By F. R. HOGG, M.D., Surgeon-Major. Crown 8vo, 5s.

Endemic Diseases of Tropical Climates, with their Treatment. By JOHN SULLIVAN, M.D. Post 8vo, 6s.

The National Dispensatory : Containing the Natural History, Chemistry, Pharmacy, Actions and Uses of Medicines. By ALFRED STILLÉ, M.D., LL.D., and JOHN M. MAISCH, Ph.D. Second Edition. 8vo, with 239 Engravings, 34s.

Binz's Elements of Therapeutics : A Clinical Guide to the Action of Drugs. Translated by E. I. SPARKS, M.B., F.R.C.P. Crown 8vo, 8s. 6d.

The Pharmacopœia of the London Hospital. Compiled under the direction of a Committee appointed by the Hospital Medical Council. Fcp. 8vo, 3s.

A Companion to the British Pharmacopœia. By PETER SQUIRE, F.L.S., assisted by his sons, P. W. and A. H. SQUIRE. Thirteenth Edition. 8vo, 10s. 6d.

By the same Authors.

The Pharmacopœias of the London Hospitals, arranged in Groups for Easy Reference and Comparison. Fourth Edition. 18mo, 6s.

Bazaar Medicines of India, And Common Medical Plants: With Full Index of Diseases, indicating their Treatment by these and other Agents procurable throughout India, &c. By E. J. WARING, C.I.E., M.D., F.R.C.P. Third Edition. Fcap. 8vo, 5s.

Tropical Dysentery and Chronic Diarrhœa—Liver Abscess—Malarial Cachexia—Insolation—with other forms of Tropical Diseases, &c. By Sir JOSEPH FAYRER, K.C.S.I., M.D., 8vo., 15s.

By the same Author.

Climate and Fevers of India, with a series of Cases (Croonian Lectures, 1882). 8vo, with 17 Temperature Charts, 12s.

Also,

Clinical and Pathological Observations in India. 8vo, with Engravings, 20s.

Family Medicine for India. By WILLIAM J. MOORE, M.D., Honorary Surgeon to the Viceroy of India. Published under the Authority of the Government of India. Fourth Edition. Post 8vo, with Engravings. (*In the Press.*)

By the same Author.

Health-Resorts for Tropical Invalids, in India, at Home, and Abroad. Post 8vo, 5s.

The Elements of Indian Hygiene. Intended to guide the Public in acquiring some knowledge in the all-important subject of the Preservation of Health and Prevention of Sickness. By JOHN C. LUCAS, F.R.C.S., H.M.'s Indian Medical Service. Crown 8vo, with Map of India, &c., 5s.

Spirillum Fever:

(Synonyms, Famine or Relapsing Fever), as seen in Western India. By H. VANDYKE CARTER, M.D., Surgeon-Major I.M.D. 8vo, with Plates, 21s.

Diseases of Tropical Climates,

And their Treatment: with Hints for the Preservation of Health in the Tropics. By JAMES A. HORTON, M.D., Surgeon-Major. Second Edition. Post 8vo, 12s. 6d.

The Student's Guide to the

Practice of Medicine. By MATTHEW CHARTERIS, M.D., Professor of Materia Medica in the University of Glasgow. Third Edition. Fcap. 8vo, with Engravings on Copper and Wood, 7s.

Hooper's Physicians' Vade-

Mecum. A Manual of the Principles and Practice of Physic. Tenth Edition. By W. A. GUY, F.R.C.P., F.R.S., and J. HARLEY, M.D., F.R.C.P. With 118 Engravings. Fcap. 8vo, 12s. 6d.

Clinical Studies:

Illustrated by Cases observed in Hospital and Private Practice. By Sir J. ROSE CORMACK, M.D., F.R.S.E., late Physician to the Hertford British Hospital of Paris. Two Vols. Post 8vo, 20s.

Clinical Medicine:

Lectures and Essays. By BALTHAZAR FOSTER, M.D., F.R.C.P. Lond., Professor of Medicine in Queen's College, Birmingham. 8vo, 10s. 6d.

Clinical Lectures and Cases,

with Commentaries. By HENRY THOMPSON, M.D., F.R.C.P., Consulting Physician to Middlesex Hospital. With Temperature Charts. 8vo, 7s. 6d.

Clinical Medicine:

A Systematic Treatise on the Diagnosis and Treatment of Disease. By AUSTIN FLINT, M.D., Professor of Medicine in the Bellevue Hospital Medical College. 8vo, 20s.

By the same Author.

Phthisis:

In a series of Clinical Studies. 8vo, 16s.

Transfusion of Human Blood:

With Table of 50 cases. By Dr. ROUSSEL, of Geneva. With a Preface by Sir JAMES PAGET, Bart. Crown 8vo, 2s. 6d.

The Student's Guide to Medical

Case-Taking. By FRANCIS WARNER, M.D., Assistant Physician to the London Hospital. Fcap. 8vo, 5s.

How to Examine the Chest:

Being a practical Guide for the use of Students. By SAMUEL WEST, M.D., Physician to the City of London Hospital for Diseases of the Chest; Medical Tutor and Registrar at St. Bartholomew's Hospital. With numerous Engravings. Fcap. 8vo.

The Microscope in Medicine.

By LIONEL S. BEALE, M.B., F.R.S., Physician to King's College Hospital. Fourth Edition. 8vo, with 86 Plates, 21s.

Also.

On Slight Ailments:

Their Nature and Treatment. Second Edition. 8vo, 5s.

The Spectroscope in Medicine.

By CHARLES A. MACMUNN, B.A., M.D. 8vo, with 3 Chromo-lithographic Plates of Physiological and Pathological Spectra, and 13 Engravings, 9s.

The Student's Guide to Medical

Diagnosis. By SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London Hospital. Fifth Edition. Fcap. 8vo, with 111 Engravings, 7s.

By the same Author.

The Student's Outlines of Medi-

cal Treatment. Second Edition. Fcap. 8vo, 7s.

Also.

On Chronic Atrophy of the

Stomach, and on the Nervous Affections of the Digestive Organs. 8vo, 8s.

Notes on Asthma:

Its Forms and Treatment. By JOHN C. THOROWOOD, M.D., Physician to the Hospital for Diseases of the Chest. Third Edition. Crown 8vo, 4s. 6d.

Observations on the Result of

Treatment of nearly One Hundred Cases of Asthma. By T. L. PRIDHAM, M.R.C.S. Third Edition. 8vo, 2s. 6d.

The Contagiousness of Pulmo-

nary Consumption and its Anti-septic Treatment. By J. BURNEY YEO, M.D., Physician to King's College Hospital. Crown 8vo, 3s. 6d.

Diseases of the Chest:

Contributions to their Clinical History, Pathology, and Treatment. By A. T. HOUGHTON WATERS, M.D., Physician to the Liverpool Royal Infirmary. Second Edition. 8vo, with Plates, 15s.

Winter Cough:

(Catarrh, Bronchitis, Emphysema, Asthma). By HORACE DOBELL, M.D., Consulting Physician to the Royal Hospital for Diseases of the Chest. Third Edition. 8vo, with Coloured Plates, 10s. 6d.

By the same Author.

Loss of Weight, Blood-Spit-

ting, and Lung Disease. Second Edition, to which is added Part VI., "On the Functions and Diseases of the Liver." 8vo, with Chromo-lithograph, 10s. 6d.

Also.

The Mont Dore Cure, and the Proper Way to Use it. 8vo, 7s. 6d.

- Croonian Lectures on Some Points in the Pathology and Treatment of Typhoid Fever.** By WILLIAM CAYLEY, M.D., F.R.C.P., Physician to the Middlesex and the London Fever Hospitals. Crown 8vo, 4s. 6d.
- Relapse of Typhoid Fever, especially with reference to the Temperature.** By J. PEARSON IRVINE, M.D., F.R.C.P., late Assistant Physician to Charing Cross Hospital. 8vo, with Engravings, 6s.
- Diseases of the Heart and Aorta:** Clinical Lectures. By GEORGE W. BALFOUR, M.D., F.R.C.P., and F.R.S. Edin., late Senior Physician to, and Lecturer on Clinical Medicine in, the Royal Infirmary, Edinburgh. Second Edition. 8vo, with Chromo-Lithograph and Wood Engravings, 12s. 6d.
- On Diseases of the Heart.** By T. B. PEACOCK, M.D., F.R.C.P. (1) Malformations. 8vo, 10s. (2) Causes and Effects of Valvular Disease. 8vo, 5s. (3) Prognosis in Valvular Disease. 8vo, 3s. 6d.
- Manual of the Physical Diagnosis of Diseases of the Heart, including the use of the Sphygmograph and Cardiograph.** By A. E. SANSOM, M.D., F.R.C.P., Assistant-Physician to the London Hospital. Third Edition. Fcap. 8vo, with 48 Engravings, 7s. 6d.
By the same Author.
- The Antiseptic System in Medicine and Surgery: A Treatise on Carbolic Acid and its Compounds, etc.** With 9 Plates (42 Figures), 8vo, 10s. 6d.
- Medical Ophthalmoscopy:** A Manual and Atlas. By WILLIAM R. GOWERS, M.D., F.R.C.P., Assistant Professor of Clinical Medicine in University College, and Senior Assistant-Physician to the Hospital. Second Edition, with Coloured Autotype and Lithographic Plates and Woodcuts. 8vo, 18s.
By the same Author.
- Epilepsy, and other Chronic Convulsive Diseases: Their Causes, Symptoms, and Treatment.** 8vo, 10s. 6d.
Also.
- Pseudo-Hypertrophic Muscular Paralysis: A Clinical Lecture.** 8vo, with Engravings and Plate, 3s. 6d.
Also.
- The Diagnosis of Diseases of the Spinal Cord.** Second Edition. 8vo, with Coloured Plate and Engravings, 4s. 6d.
- Studies on Functional Nervous Disorders.** By C. HANDFIELD JONES, M.B., F.R.S., Physician to St. Mary's Hospital. Second Edition. 8vo, 18s.
- Diseases of the Liver:** With and without Jaundice. By GEORGE HARLEY, M.D., F.R.C.P., F.R.S. 8vo, with 2 Plates and 36 Engravings, 21s.
- Diseases of the Stomach:** The Varieties of Dyspepsia, their Diagnosis and Treatment. By S. O. HABERSHON, M.D., F.R.C.P., late Senior Physician to Guy's Hospital. Third Edition. Crown 8vo, 5s.
By the same Author.
- Pathology of the Pneumogastric Nerve, being the Lumleian Lectures for 1876.** Post 8vo, 3s. 6d.
Also.
- Diseases of the Abdomen,** Comprising those of the Stomach and other parts of the Alimentary Canal, Œsophagus, Cæcum, Intestines, and Peritoneum. Third Edition. 8vo, with 5 Plates, 21s.
- Gout, Rheumatism,** And the Allied Affections; with a Chapter on Longevity and the Causes Antagonistic to it. By PETER HOOD, M.D. Second Edition. Crown 8vo, 10s. 6d.
- Notes on Rheumatism.** By JULIUS POLLOCK, M.D., F.R.C.P., Senior Physician to the Charing Cross Hospital. Second Edition. Fcap. 8vo, with Engravings, 3s. 6d.
- Diseases of the Nervous System:** Clinical Lectures. By THOMAS BUZZARD, M.D., F.R.C.P., Physician to the National Hospital for the Paralysed and Epileptic. With Engravings, 8vo, 15s.
- A Treatise on the Diseases of the Nervous System.** By JAMES ROSS, M.D., F.R.C.P., Assistant-Physician to the Manchester Royal Infirmary. Two Vols., 8vo, with Lithographs, Photographs, and 280 Wood Engravings, 42s.
- Nervous Diseases:** Their Description and Treatment. A Manual for Students and Practitioners of Medicine. By ALLEN M. HAMILTON, M.D., Physician at the Epileptic and Paralytic Hospital, New York. Second Edition. Royal 8vo, with 72 Engravings, 16s.
- Nerve Vibration and Excitation, as Agents in the Treatment of Functional Disorder and Organic Disease.** By J. MORTIMER GRANVILLE, M.D. 8vo, 5s.
- Fits:** Diagnosis and Immediate Treatment of Cases of Insensibility and Convulsions. By JOHN H. WATERS, M.D., K.C., St.G.C., Surgeon to the C Division of Metropolitan Police. Crown 8vo, bound in leather, 4s.
- Imperfect Digestion:** Its Causes and Treatment. By A. LEARED, M.D., Seventh Edition. Fcap. 8vo, 4s. 6d.

Headaches :

Their Nature, Causes, and Treatment. By WILLIAM H. DAY, M.D., Physician to the Samaritan Hospital for Women and Children. Third Edition. Crown 8vo, with Engravings, 6s. 6d.

On Megrin, Sick Headache and some Allied Disorders : a Contribution to the Pathology of Nerve Storms. By E. LIVEING, M.D., F.R.C.P. 8vo, 15s.

Nutrition in Health and Disease : A Contribution to Hygiene and to Clinical Medicine. By HENRY BENNET, M.D. Third (Library) Edition. 8vo, 7s. Cheap Edition. Fcap. 8vo, 2s. 6d.

Food and Dietetics,

Physiologically and Therapeutically Considered. By F. W. PAVY, M.D., F.R.S., Physician to Guy's Hospital. Second Edition. 8vo, 15s.

By the same Author.

Croonian Lectures on Certain Points connected with Diabetes. 8vo, 4s. 6d.

Indigestion :

What it is; what it leads to; and a New Method of Treating it. By JOHN B. GILL, M.D. Second Edition. Fcap. 8vo, 4s. 6d.

The Climate of the Undercliff, Isle of Wight, as deduced from forty years' consecutive Meteorological Observations. By J. L. WHITEHEAD, M.D. Royal 8vo, 5s.

The Riviera :

Sketches of the Health-Resorts of the North Mediterranean Coast of France and Italy, from Hyères to Spezia; its Medical Aspect and Value, &c. By EDWARD I. SPARKS, M.B., F.R.C.P. Crown 8vo, 8s. 6d.

Winter and Spring

On the Shores of the Mediterranean. By HENRY BENNET, M.D. Fifth Edition. Post 8vo, with numerous Plates, Maps, and Engravings, 12s. 6d.

By the same Author.

Treatment of Pulmonary Consumption by Hygiene, Climate, and Medicine. Third Edition. 8vo, 7s. 6d.

The Principal Southern and Swiss Health-Resorts : their Climate and Medical Aspect. By WILLIAM MARCET, M.D., F.R.C.P., F.R.S. With Illustrations. Crown 8vo, 7s. 6d.

The Ocean as a Health-Resort : A Practical Handbook of the Sea, for the use of Tourists and Health-Seekers. By WILLIAM S. WILSON, L.R.C.P., Second Edition, with Chart of Ocean Routes, &c. Crown 8vo, 7s. 6d.

Davos Platz, and the Effects of High Altitude on Phthisis. By ALFRED WISE, M.D. Fcap. 8vo, 2s. 6d.

Principal Health-Resorts

Of Europe and Africa, and their Use in the Treatment of Chronic Diseases. By THOMAS MORE MADDEN, M.D., M.R.I.A. 8vo, 10s.

Handbook of Medical and Surgical Electricity. By HERBERT TIBBITS, M.D., F.R.C.P.E., Senior Physician to the West London Hospital for Paralysis and Epilepsy. Second Edition. 8vo, with 95 Engravings, 9s.

By the same Author.

A Map of Ziemssen's Motor Points of the Human Body : A Guide to Localised Electrification. Mounted on Rollers, 35 x 21. With 20 Illustrations, 5s.

Lectures on the Clinical Uses of Electricity. By J. RUSSELL REYNOLDS, M.D., F.R.S., Physician to University College Hospital. Second Edition. Post 8vo, 3s. 6d.

A System of Practical Surgery. By Sir WILLIAM FERGUSSON, Bart., F.R.S. Fifth Edition. 8vo, with 463 Engravings, 21s.

Surgical Emergencies :

Together with the Emergencies Attendant on Parturition and the Treatment of Poisoning. By PAUL SWAIN, F.R.C.S., Surgeon to the Royal Albert Hospital, Devonport. Third Edition. Crown 8vo, with 117 Engravings, 5s.

A Course of Operative Surgery. By CHRISTOPHER HEATH, Surgeon to University College Hospital. With 20 Plates drawn from Nature by M. LÉVEILLÉ, and coloured by hand under his direction. Large 8vo, 40s.

By the same Author.

The Student's Guide to Surgical Diagnosis. Fcap. 8vo, 6s. 6d.

Also.

Manual of Minor Surgery and Bandaging. For the use of House Surgeons, Dressers, and Junior Practitioners. Sixth Edition. Fcap. 8vo, with 115 Engravings, 5s. 6d.

Also.

Injuries and Diseases of the Jaws. Second Edition. 8vo, with 164 Engravings, 12s.

Outlines of Surgery and Surgical Pathology. By F. LE GROS CLARK, F.R.S., assisted by W. W. WAGSTAFFE, F.R.C.S. Second Edition. 8vo, 10s. 6d.

Regional Surgery :

Including Surgical Diagnosis. A Manual for the use of Students. Part I. The Head and Neck. By F. A. SOUTHAM, M.A., M.B., F.R.C.S., Assistant-Surgeon to the Manchester Royal Infirmary. Crown 8vo, 6s. 6d.

The Practice of Surgery :

A Manual. By THOMAS BRYANT, Surgeon to Guy's Hospital. Third Edition. Two Vols. Crown 8vo, with 672 Engravings (many being coloured), 28s.

The Surgeon's Vade-Mecum :

A Manual of Modern Surgery. By ROBERT DRUITT, F.R.C.S. Eleventh Edition. Fcap. 8vo, with 369 Engravings, 14s.

Illustrations of Clinical Surgery.

By JONATHAN HUTCHINSON, Senior Surgeon to the London Hospital. In occasional fasciculi. I. to XV., 6s. 6d. each. Fasciculi I. to X. bound, with Appendix and Index, £3 10s.

The Principles and Practice of Surgery.

By WILLIAM PIRRIE, F.R.S.E., late Professor of Surgery in the University of Aberdeen. Third Edition. 8vo, with 490 Engravings, 28s.

Surgical Enquiries :

Including the Hastings Essay on Shock, the Treatment of Inflammations, and numerous Clinical Lectures. By FURNEAUX JORDAN, F.R.C.S., Professor of Surgery, Queen's College, Birmingham. Second Edition, with numerous Plates. Royal 8vo, 12s. 6d.

Treatment of Wounds :

Clinical Lectures. By SAMPSON GAMGEE, F.R.S.E., Surgeon to the Queen's Hospital, Birmingham. Crown 8vo, with Engravings, 5s.

By the same Author.

Fractures of the Limbs,

And their Treatment. 8vo, with Plates, 10s. 6d.

On Dislocations and Fractures.

By JOSEPH MACLISE, F.R.C.S. Uniform with "Surgical Anatomy." 36 folio Plates and Text. Cloth, £2 10s.

Injuries of the Spine and Spinal

Cord, without Apparent Mechanical Lesion, and NERVOUS SHOCK, in their Surgical and Medico-Legal Aspects. By HERBERT W. PAGE, M.A., M.C. Cantab., F.R.C.S., Surgeon to St. Mary's Hospital. 8vo, 12s. 6d.

Lectures on Orthopædic Surgery.

By BERNARD E. BRODHURST, F.R.C.S., Surgeon to the Royal Orthopædic Hospital. Second Edition. 8vo, with Engravings, 12s. 6d.

By the same Author.

On Anchylosis, and the Treat-

ment for the Removal of Deformity and the Restoration of Mobility in Various Joints. Fourth Edition. 8vo, with Engravings, 5s.

Orthopædic Surgery,

And Diseases of the Joints. By L. A. SAYRE, M.D., Professor of Orthopædic Surgery in Bellevue Hospital Medical College. 8vo, with 274 Engravings, 20s.

Clubfoot :

Its Causes, Pathology, and Treatment. By WM. ADAMS, F.R.C.S., Surgeon to the Great Northern Hospital. Second Edition. 8vo, with 106 Engravings and 6 Lithographic Plates, 15s.

By the same Author.

On Contraction of the Fingers,

And its Treatment by Subcutaneous Operation; and on Obliteration of Depressed Cicatrices, by the same Method. 8vo, with 30 Engravings, 4s. 6d.

Also.

Lateral and other Forms of Curvature of the Spine: Their Pathology and Treatment. Second Edition. 8vo, with 5 Lithograph Plates and 72 Wood Engravings, 10s. 6d.**Osteotomy :**

With an Enquiry into the Etiology and Pathology of Knock-knee, Bow-leg, and other Osseous Deformities of the Lower Limbs. By WILLIAM MACEWEN, M.D., Surgeon and Lecturer on Clinical Surgery to the Glasgow Royal Infirmary. 8vo, with 51 Engravings, 7s. 6d.

Lectures on Diseases of Bones and Joints.

By CHARLES MACNAMARA, F.R.C.S., Surgeon to, and Lecturer on Surgery at, Westminster Hospital. Crown 8vo, with Engravings, 10s. 6d.

Orthopraxy :

The Mechanical Treatment of Deformities, Debilities, and Deficiencies of the Human Frame. By H. HEATHER BIGG, Assoc. Inst. C.E. Third Edition. 8vo, with 319 Engravings, 15s.

The Orthopragms of the Spine :

An Essay on the Curative Mechanisms applicable to Spinal Curvature, etc. By ROBERT HEATHER BIGG, Assoc. Inst. C.E. 8vo, with Engravings, 5s.

On Diseases and Injuries of the

Eye: A Course of Systematic and Clinical Lectures to Students and Medical Practitioners. By J. R. WOLFE, M.D., F.R.C.S.E., Senior Surgeon to the Glasgow Ophthalmic Institution; Lecturer on Ophthalmic Medicine and Surgery in Anderson's College. With 10 Coloured Plates and 157 Wood Engravings. 8vo, £1 1s.

Hints on Ophthalmic Out-Patient

Practice. By CHARLES HIGGENS, Ophthalmic Surgeon to, and Lecturer on Ophthalmology at, Guy's Hospital. Second Edition. Fcap. 8vo, 3s.

Liebreich's Atlas of Ophthal-

moscopy: Composed of 12 Chromolithographic Plates (containing 59 Figures). The Text translated by H. ROSBOROUGH SWANZY, M.B. Second Edition. 4to, 30s.

The Student's Guide to Diseases of the Eye. By EDWARD NETTLESHIP, F.R.C.S., Ophthalmic Surgeon to St. Thomas's Hospital and to the Hospital for Sick Children. Second Edition. Fcap. 8vo, with 137 Engravings, 7s. 6d.

A Manual of Diseases of the Eye. By C. MACNAMARA, F.R.C.S., Surgeon to Westminster Hospital. Fourth Edition. Crown 8vo, with 4 Coloured Plates and 66 Engravings, 10s. 6d.

A Manual of the Principles and Practice of Ophthalmic Medicine and Surgery. By T. WHARTON JONES, F.R.C.S., F.R.S. Third Edition. Fcap. 8vo, with 9 Coloured Plates and 173 Engravings, 12s. 6d.

Glaucoma :

Its Causes, Symptoms, Pathology, and Treatment. By PRIESTLEY SMITH, M.R.C.S., Ophthalmic Surgeon to the Queen's Hospital, Birmingham. 8vo, with Lithographic Plates, 10s. 6d.

A Manual of Ophthalmoscopy for the use of Students. By DR. DAGUENET. Translated by C. S. JEAFFRESON, Surgeon to the Newcastle-on-Tyne Eye Infirmary. With Engravings. Fcap. 8vo, 5s.

Essays in Ophthalmology.

By GEORGE E. WALKER, F.R.C.S., Surgeon to St. Paul's Eye and Ear Hospital, &c., Liverpool. Post 8vo, 6s.

Hare-Lip and Cleft Palate.

By FRANCIS MASON, F.R.C.S., Surgeon to, and Lecturer on Practical Surgery at, St. Thomas's Hospital. 8vo, with 66 Engravings, 6s.

By the same Author.

The Surgery of the Face.

8vo, with 100 Engravings, 7s. 6d.

A Practical Treatise on Aural Surgery. By H. MACNAUGHTON JONES, M.D., Professor of the Queen's University in Ireland, Surgeon to the Cork Ophthalmic and Aural Hospital. Second Edition. Crown 8vo, with 63 Engravings, 8s. 6d.

By the same Author.

Atlas of Diseases of the Membrana Tympani. In Coloured Plates, containing 62 Figures, with Text. Crown 4to, 21s.

Diseases and Injuries of the Ear. By WILLIAM B. DALBY, F.R.C.S., Aural Surgeon to, and Lecturer on Surgery at, St. George's Hospital. Second Edition. Fcap. 8vo, with Engravings, 6s. 6d.

Lectures on Syphilis of the Larynx (Lesions of the Secondary and Intermediate Stages). By W. MACNEILL WHISTLER, M.D., Physician to the Hospital for Diseases of the Throat and Chest. Post 8vo, 4s.

Diseases of the Throat and Nose. A Manual. By MORELL MACKENZIE, M.D. Lond., Senior Physician to the Hospital for Diseases of the Throat and Chest. Vol. I. Diseases of the Pharynx, Larynx, and Trachea. Post 8vo, with 112 Engravings, 12s. 6d.

By the same Author.

Diphtheria :

Its Nature and Treatment, Varieties, and Local Expressions. 8vo, 5s.

Sore Throat :

Its Nature, Varieties, and Treatment. By PROSSER JAMES, M.D., Physician to the Hospital for Diseases of the Throat. Fourth Edition. Post 8vo, with Coloured Plates and Engravings, 6s. 6d.

Throat Diseases,

And the Use of the Laryngoscope: A Handbook for Practitioners and Senior Students. By W. D. HEMMING, F.R.C.S.E. With Engravings. Fcap. 8vo, 2s. 6d.

The Ear :

Its Anatomy, Physiology, and Diseases. By C. H. BURNETT, A.M., M.D., Aural Surgeon to the Presbyterian Hospital, Philadelphia. 8vo, with 87 Engravings, 18s.

A Treatise on Vocal Physiology and Hygiene, with especial reference to the Cultivation and Preservation of the Voice. By GORDON HOLMES, M.D., Physician to the Municipal Throat and Ear Infirmary. Second Edition. With Engravings. Crown 8vo, 6s. 6d.

By the same Author.

A Guide to the Use of the Laryngoscope in General Practice. Crown 8vo, with Engravings, 2s. 6d.

A System of Dental Surgery.

By JOHN TOMES, F.R.S., and C. S. TOMES, M.A., F.R.S. Second Edition. Fcap. 8vo, with 268 Engravings, 14s.

Dental Anatomy, Human and Comparative: a Manual. By CHARLES S. TOMES, M.A., F.R.S. Second Edition. Crown 8vo, with 191 Engravings, 12s. 6d.

A Practical Treatise on Operative Dentistry. By JONATHAN TAFT, D.D.S., Professor in the Ohio College of Dental Surgery. Third Edition. With 134 Engravings. 8vo, 18s.

The Student's Guide to Dental Anatomy and Surgery. By HENRY SEWILL, M.R.C.S., L.D.S. Second Edition. Fcap. 8vo, with Engravings.

(In the Press.)

Elements of Dental Materia Medica and Therapeutics, with Pharmacopœia. By JAMES STOCKEN, L.D.S.R.C.S., Pereira Prizeman for Materia Medica, and THOMAS GADDES, L.D.S. Eng. and Edin. Third Edition. Fcap. 8vo, 7s. 6d.

A Manual of Dental Mechanics.

By OAKLEY COLES, L.D.S.R.C.S.,
Second Edition. Crown 8vo, with 140
Engravings, 7s. 6d.

By the same Author.

Deformities of the Mouth.

Third Edition, 8vo, with 83 Wood En-
gravings and 96 Drawings on Stone, 12s. 6d.

Mechanical Dentistry in Gold and Vulcanite.

By F. H. BALK-
WILL, L.D.S.R.C.S. 8vo, with 2 Litho-
graphic Plates and 57 Engravings, 10s.

Lectures on Dermatology :

Delivered at the Royal College of Sur-
geons, by Sir ERASMUS WILSON, F.R.S.
1870, 6s. ; 1871-73, 10s. 6d. ; 1874-75,
10s. 6d. ; 1876-78, 10s. 6d.

Eczema :

By MCCALL ANDERSON, M.D., Professor
of Clinical Medicine in the University of
Glasgow. Third Edition. 8vo, with
Engravings, 7s. 6d.

Eczema and its Management :

A practical Treatise based on the Study
of 2,500 Cases of the Disease. By L.
DUNCAN BULKLEY, M.D., Physician for
Skin and Venereal Diseases at the New
York Hospital. 8vo, 12s. 6d.

By the same Author.

Diseases of the Skin :

A Manual, with an Analysis of 8,000 Con-
secutive Cases and a Formulary. Crown
8vo, 6s. 6d.

Psoriasis, or Lepra.

By GEORGE GASKOIN, M.R.C.S., Sur-
geon to the British Hospital for Diseases
of the Skin. 8vo, 5s.

On Certain Rare Diseases of the

Skin : being Vol. 1 of "Lectures on
Clinical Surgery." By JONATHAN
HUTCHINSON, Senior Surgeon to the
London Hospital, and to the Hospital
for Diseases of the Skin. 8vo, 10s. 6d.

Leprosy in British Guiana :

An Account of West Indian Leprosy. By
JOHN D. HILLIS, F.R.C.S., M.R.I.A.,
Medical Superintendent of the Leper
Asylum, British Guiana. Imp. 8vo, with
22 Lithographic Coloured Plates and
Wood Engravings. £1 11s. 6d.

Photographic Illustrations of

Skin Diseases. Sixty Cases from Life.
By GEORGE H. FOX, M.D. 4to, £5 5s.

Atlas of Skin Diseases :

By TILBURY FOX, M.D., F.R.C.P.
With 72 Coloured Plates. Royal 4to, half
morocco, £6 6s.

Sarcoma and Carcinoma :

Their Pathology, Diagnosis, and Treat-
ment. By HENRY T. BUTLIN, F.R.C.S.,
Assistant-Surgeon to St. Bartholomew's
Hospital. 8vo, with 4 Plates, 8s.

Cancer of the Breast :

By THOMAS W. NUNN, F.R.C.S., Con-
sulting Surgeon to the Middlesex Hos-
pital. 4to, with 21 Coloured Plates, £2 2s.

Cancer Life :

Its Causes, Progress, and Treatment. - A
General and Historical Treatise. By
R. MITCHELL, M.R.C.S. 8vo, 7s. 6d.

Certain Forms of Cancer,

With a New and Successful Mode of treat-
ing it. By A. MARSDEN, Senior Surgeon
to the Cancer Hospital. Second Edition.
8vo, with Coloured Plates, 8s. 6d.

On Cancer :

Its Allies, and other Tumours, with special
reference to their Medical and Surgical
Treatment. By F. A. PURCELL, M.D.,
M.C., Surgeon to the Cancer Hospital,
Brompton. 8vo, with 21 Engravings,
10s. 6d.

Diseases of the Urinary Organs :

Clinical Lectures. By Sir HENRY
THOMPSON, F.R.C.S., Emeritus Pro-
fessor of Clinical Surgery in University
College. Sixth (Students') Edition. 8vo,
with 73 Engravings, 2s. 6d.

By the same Author.

Diseases of the Prostate :

Their Pathology and Treatment. Fifth
(Students') Edition. 8vo, with numerous
Engravings, 2s. 6d.

Also.

Practical Lithotomy and Litho-

trity ; or, an Inquiry into the best Modes
of Removing Stone from the Bladder.
Third Edition. 8vo, with 87 Engravings,
10s.

Also.

The Preventive Treatment of

Calculous Disease, and the Use of
Solvent Remedies. Second Edition.
Fcap. 8vo, 2s. 6d.

Diseases of the Testis, Sperm-

atic Cord, and Scrotum. By
THOMAS B. CURLING, F.R.S., Consult-
ing Surgeon to the London Hospital.
Fourth Edition. 8vo, with Engravings, 16s.

Fistula, Hæmorrhoids, Painful

**Ulcer, Stricture, Prolapsus, and
other Diseases of the Rectum :**
Their Diagnosis and Treatment. By
WILLIAM ALLINGHAM, Surgeon to St.
Mark's Hospital for Fistula. Fourth
Edition. 8vo, with Engravings, 10s. 6d.

Hæmorrhoidal Disorder.

By JOHN GAY, F.R.C.S., Senior Sur-
geon to the Great Northern Hospital.
8vo, with Engravings, 2s. 6d.

Hydrocele :

Its several Varieties and their Treatment.
By SAMUEL OSBORN, late Surgical
Registrar to St. Thomas's Hospital.
Fcap. 8vo, with Engravings, 3s.

By the same Author.

Diseases of the Testis.

Fcap. 8vo, with Engravings, 3s. 6d.

Parasites :

A Treatise on the Entozoa of Man and Animals, including some Account of the Ectozoa. By T. SPENCER COBBOLD, M.D., F.R.S. 8vo, with 85 Engravings, 15s.

The Surgery of the Rectum.

By HENRY SMITH, Professor of Surgery in King's College, Surgeon to the Hospital. Fifth Edition. 8vo, 6s.

Cancer of the Rectum :

Its Pathology, Diagnosis, and Treatment. By W. HARRISON CRIPPS, F.R.C.S., Assist.-Surg. to St. Bartholomew's Hospital, &c. Cr. 8vo, with Lithographic Plates, 6s.

Lectures on the Surgical Disorders of the Urinary Organs. By REGINALD HARRISON, F.R.C.S., Surgeon to the Liverpool Royal Infirmary. Second Edition, with 48 Engravings. 8vo, 12s. 6d.

By the same Author.

The Prevention of Stricture and of Prostatic Obstruction. 8vo, with Engravings, 2s. 6d.

Lithotomy and Extraction of Stone. By W. P. HARRIS, M.D., Surgeon-Major H.M. Bengal Medical Service. 8vo, with Engravings, 10s. 6d.

Diseases of the Bladder,

Prostate Gland, and Urethra, with a practical view of Urinary Diseases, Deposits, and Calculi. By F. J. GANT, Senior Surgeon to the Royal Free Hospital. Fourth Edition. Crown 8vo, 10s. 6d.

Morbid Conditions of the Urine, dependent upon Derangements of Digestion. By CHARLES H. RALFE, M.D., F.R.C.P., Assistant-Physician to the London Hospital. Crown 8vo, 6s.

Renal and Urinary Diseases :

Clinical Reports. By WILLIAM CARTER, M.B., Physician to the Liverpool Southern Hospital. Crown 8vo, 7s. 6d.

Pathology of the Urine,

Including a Complete Guide to its Analysis. By J. L. W. THUDICHUM, M.D., F.R.C.P. Second Edition, rewritten and enlarged. 8vo, with Engravings, 15s.

Genito-Urinary Organs, including Syphilis : a Practical Treatise on their Surgical Diseases, designed as a Manual for Students and Practitioners. By W. H. VAN BUREN, M.D., and E. L. KEYES, M.D. Royal 8vo, with 140 Engravings, 21s.

Lectures on Syphilis.

By HENRY LEE, Consulting Surgeon to St. George's Hospital. 8vo, 10s.

Harveian Lectures on Syphilis.

By JAMES R. LANE, F.R.C.S., late Surgeon to St. Mary's Hospital. Second Edition. Fcap. 8vo, 3s. 6d.

Photographic Illustrations of Cutaneous Syphilis. Seventy Cases from Life. By G. H. FOX, M.D. 4to, £5 5s.

Urinary and Reproductive Organs : their Functional Diseases. By D. CAMPBELL BLACK, M.D. Second Edition. 8vo, 10s.

A Treatise on Syphilis.

By WALTER J. COULSON, Surgeon to the Lock Hospital and to St. Peter's Hospital for Stone. 8vo, 10s.

By the same Author.

Stone in the Bladder :

Its Prevention, early Symptoms, and Treatment by Lithotripsy. 8vo, 6s.

Also.

Coulson on Diseases of the Bladder and Prostate Gland. Sixth Edition. 8vo, 16s.

The Reproductive Organs

In Childhood, Youth, Adult Age, and Advanced Life, considered in their Physiological, Social, and Moral Relations. By WILLIAM ACTON, M.R.C.S. Sixth Edition. 8vo, 12s.

Student's Primer on the Urine.

By J. TRAVIS WHITTAKER, M.D., Clinical Demonstrator at the Royal Infirmary, Glasgow. With 16 Plates etched on Copper. Post 8vo, 4s. 6d.

A Manual of the Laws affecting Medical Men. By ROBERT G. GLENN, LL.B., Barrister-at-Law. 8vo, 14s.

The Medical Adviser in Life Assurance. By EDWARD H. SIEVEKING, M.D., F.R.C.P., Physician to St. Mary's and Lock Hospitals, &c. Second Edition. Crown 8vo, 6s.

A Dictionary of Medical Science : Containing a concise Explanation of the various Subjects and Terms of Medicine, &c. ; Notices of Climate and Mineral Waters ; Formulæ for Official, Empirical, and Dietetic Preparations ; with the Accentuation and Etymology of the Terms, and the French and other Synonyms. By ROBLEY DUNGLISON, M.D., LL.D. New Edition. Royal 8vo, 28s.

A Medical Vocabulary :

Being an Explanation of all Terms and Phrases used in the various Departments of Medical Science and Practice, giving their Derivation, Meaning, Application, and Pronunciation. By ROBERT G. MAYNE, M.D., LL.D. Fifth Edition. Fcap. 8vo, 10s. 6d.

Abridged Medical Account Books. The "Expedite" Method. By JAMES MACNAB, L.R.C.S.E. *Index Ledger.* Royal 4to. For three years, 15s. *Visiting List.* Cloth, 2s. 6d. ; Leather, 3s. 6d.

Medical Education

And Practice in all parts of the World. By HERBERT JUNIUS HARDWICKE, M.D., M.R.C.P. 8vo, 10s.

INDEX.

- Acton's Reproductive Organs, 14
 Adams (W.) on Clubfoot, 11
 ——— Contraction of the Fingers, 11
 ——— Curvature of the Spine, 11
 Allan on Fever Nursing, 7
 Allingham on Diseases of the Rectum, 13
 Anatomical Remembrancer, 4
 Anderson (McC.) on Eczema, 13
 Aveling on the Chamberlens and Midwifery Forceps, 6
 ——— on the Influence of Posture on Women, 6
 Balfour's Diseases of the Heart and Aorta, 9
 Balkwill's Mechanical Dentistry, 13
 Bantock on Rupture of the Female Perineum, 6
 Barnes on Obstetric Operations, 5
 ——— on Diseases of Women, 5
 Beale's Microscope in Medicine, 8
 ——— Slight Ailments, 8
 Bellamy's Surgical Anatomy, 3
 Bennet (J. H.) on the Mediterranean, 10
 ——— on Pulmonary Consumption, 10
 ——— on Nutrition, 10
 Bentley and Trimen's Medicinal Plants, 7
 Bentley's Manual of Botany, 7
 Bigg (H. H.) on Orthopraxy, 11
 Bigg (R. H.) on the Orthopragms of Spine, 11
 ——— on Spinal Curvature, 8
 Binz's Elements of Therapeutics, 7
 Black on the Urinary Organs, 14
 Braune's Topographical Anatomy, 3
 Brodhurst's Anchylosis, 11
 ——— Orthopædic Surgery, 11
 Bryant's Practice of Surgery, 11
 Bucknill and Tuke's Psychological Medicine, 5
 Bulkley on Diseases of the Skin, 13
 ——— on Eczema, 13
 Burdett's Cottage Hospitals, 5
 ——— Pay Hospitals, 5
 Burnett on the Ear, 12
 Burton's Midwifery for Midwives, 5
 Butlin's Sarcoma and Carcinoma, 13
 Buzzard's Diseases of the Nervous System, 9
 Carpenter's Human Physiology, 4
 Carter (H. V.) on Spirillum Fever, 8
 Carter (W.) on Renal and Urinary Diseases, 14
 Cayley's Typhoid Fever, 9
 Charteris' Practice of Medicine, 8
 Clark's Outlines of Surgery, 10
 Clay's (C.) Obstetric Surgery, 6
 Cobbold on Parasites, 14
 Coles' Dental Mechanics, 13
 ——— Deformities of the Mouth, 13
 Cormack's Clinical Studies, 8
 Coulson on Diseases of the Bladder, 14
 ——— on Stone in the Bladder, 14
 ——— on Syphilis, 14
 Courty's Diseases of the Uterus, Ovaries, &c., 5
 Cripps' Cancer of the Rectum, 14
 Cullingworth's Manual of Nursing, 7
 Curling's Diseases of the Testis, 13
 Dagenet's Manual of Ophthalmoscopy, 12
 Dalby's Diseases and Injuries of the Ear, 12
 Dalton's Human Physiology, 4
 Day on Diseases of Children, 6
 ——— on Headaches, 10
 De Chaumont's Sanitary Assurance, 4
 Dobell's Lectures on Winter Cough, 8
 ——— Loss of Weight, &c., 8
 ——— Mont Dore Cure, 8
 Donville's Manual for Nurses, 7
 Druitt's Surgeon's Vade-Mecum, 11
 Duncan on the Female Perineum, 5
 ——— on Diseases of Women, 5
 Dunglison's Medical Dictionary, 14
 Ellis's Manual for Mothers, 6
 ——— of the Diseases of Children, 6
 Emmet's Gynæcology, 5
 Fayer's Climate and Fevers of India, 7
 ——— Observations in India, 7
 ——— Tropical Dysentery and Diarrhœa, 7
 Fenwick's Chronic Atrophy of the Stomach, 8
 ——— Medical Diagnosis, 8
 ——— Outlines of Medical Treatment, 8
 Fergusson's Practical Surgery, 10
 Flint on Clinical Medicine, 8
 ——— on Phthisis, 8
 Flower's Diagrams of the Nerves, 4
 Foster's Clinical Medicine, 8
 Fox's (C. B.) Examinations of Water, Air, and Food, 4
 Fox's (G. H.) Photographs of Cutaneous Syphilis, 14
 ——— Skin Diseases, 13
 Fox's (T.) Atlas of Skin Diseases, 13
 Frey's Histology and Histo-Chemistry, 4
 Fulton's Text-Book of Physiology, 4
 Galabin's Diseases of Women, 6
 Gamgee's Fractures of the Limbs, 11
 ——— Treatment of Wounds, 11
 Gant's Diseases of the Bladder, 14
 Gaskoin on Psoriasis or Lepra, 13
 Gay on Hæmorrhoidal Disorder, 13
 Gill on Indigestion, 10
 Glenn's Laws affecting Medical Men, 14
 Godlee's Atlas of Human Anatomy, 3
 Gowers' Diseases of the Spinal Cord, 9
 ——— Epilepsy, 9
 ——— Medical Ophthalmoscopy, 9
 ——— Pseudo-Hypertrophic Muscular Paralysis, 9
 Granville on Nerve Vibration and Excitation, 9
 Habershon's Diseases of the Abdomen, 9
 ——— Stomach, 9
 ——— Pneumogastric Nerve, 9
 Hamilton's Nervous Diseases, 9
 Hardwicke's Medical Education, 14
 Harley on Diseases of the Liver, 9
 Harris on Lithotomy, 14
 Harrison's Surgical Disorders of the Urinary Organs, 14
 ——— Prevention of Stricture, 14
 Heath's Injuries and Diseases of the Jaws, 10
 ——— Minor Surgery and Bandaging, 10
 ——— Operative Surgery, 10
 ——— Practical Anatomy, 3
 ——— Surgical Diagnosis, 10
 Hemming on the Laryngoscope, 12
 Higgens' Ophthalmic Out-patient Practice, 11
 Hillis' Leprosy in British Guiana, 13
 Hogg's Indian Notes, 7
 Holden's Dissections, 3
 ——— Human Osteology, 3
 ——— Landmarks, 3
 Holmes' (G.) Guide to Use of Laryngoscope, 12
 ——— Vocal Physiology and Hygiene, 12
 Hood on Gout, Rheumatism, &c., 9
 Hooper's Physicians' Vade-Mecum, 8
 Horton's Tropical Diseases, 8
 Hutchinson's Clinical Surgery, 11
 ——— Rare Diseases of the Skin, 13
 Huth's Marriage of Near Kin, 4
 Ireland's Idiocy and Imbecility, 5
 Irvine's Relapse of Typhoid Fever, 9
 James (P.) on Sore Throat, 12
 Jones' (C. H.) Functional Nervous Disorders, 9
 Jones (C. H.) and Sieveking's Pathological Anatomy, 4
 Jones' (H. McN.) Aural Surgery, 12
 ——— Atlas of Diseases of Membrana Tympani, 12
 Jones' (T. W.) Ophthalmic Medicine and Surgery, 12
 Jordan's Surgical Enquiries, 11
 Lancereaux's Atlas of Pathological Anatomy, 4
 Lane's Lectures on Syphilis, 14
 Lee (H.) on Syphilis, 14
 Leared on Imperfect Digestion, 9
 Lewis (Bevan) on the Human Brain, 4
 Liebreich's Atlas of Ophthalmoscopy, 11
 Liveing's Megrin, Sick Headache, &c., 10
 Lucas's Indian Hygiene, 7
 Macdonald's (A.) Chronic Disease of the Heart, 6
 Macdonald's (J. D.) Examination of Water, 4
 Macewen's Osteomy: Knock-knee, Bow-leg, &c., 11
 Mackenzie on Diphtheria, 12
 ——— on Diseases of the Throat and Nose, 12
 MacLise's Dislocations and Fractures, 11
 ——— Surgical Anatomy, 3
 MacMunn's Spectroscope in Medicine, 8
 Macnab's Medical Account Books, 14
 Macnamara's Diseases of Bones and Joints, 11
 ——— the Eye, 12
 Madden's Principal Health-Resorts, 10
 Marcet's Southern and Swiss Health-Resorts, 10
 Marsden's Certain Forms of Cancer, 13
 Martin's Military and State Medicine, 5
 Mason on Hare-Lip and Cleft Palate, 12
 ——— on Surgery of the Face, 12
 Mayne's Medical Vocabulary, 14
 ——— Notes on Poisons, 7
 ——— Therapeutical Remembrancer, 7
 Mitchell (R.) on Cancer Life, 13

[Continued on the next page.]

- Mitchell's (S. Weir) Nervous System in Women, 6
 Moore's Family Medicine for India, 7
 Health-Resorts for Tropical Invalids, 7
 Morris' (H.) Anatomy of the Joints, 3
 Nettleship's Diseases of the Eye, 12
 Nunn's Cancer of the Breast, 13
 Ogston's Medical Jurisprudence, 4
 Osborn on Diseases of the Testis, 13
 on Hydrocele, 13
 Page's Injuries of the Spine, 11
 Parkes' Practical Hygiene, 5
 Pavy on Diabetes, 10
 on Food and Dietetics, 10
 Peacock's Diseases of the Heart, 9
 Pharmacopœia of the London Hospital, 7
 Phillips' Materia Medica and Therapeutics, 7
 Pirrie's Principles and Practice of Surgery, 11
 Pollock on Rheumatism, 9
 Pridham on Asthma, 8
 Purcell on Cancer, 13
 Radford's Cæsarean Section, 5
 Ralfe's Morbid Conditions of the Urine, 14
 Ranisbotham's Obstetrics, 6
 Reynolds' (J. J.) Diseases of Women, 6
 Notes on Midwifery, 6
 Reynolds' (J. R.) Clinical Electricity, 10
 Roberts' (C.) Manual of Anthropometry, 5
 Roberts' (D. Lloyd) Practice of Midwifery, 5
 Ross's Diseases of the Nervous System, 9
 Roth on Dress: Its Sanitary Aspect, 5
 Roussel's Transfusion of Blood, 8
 Routh's Infant Feeding, 6
 Royle and Harley's Materia Medica, 7
 Sanderson's Physiological Handbook, 4
 Sanson's Diseases of the Heart, 9
 Antiseptic System, 9
 Savage on the Female Pelvic Organs, 6
 Sayre's Orthopædic Surgery, 11
 Schroeder's Manual of Midwifery, 6
 Sewill's Dental Anatomy, 12
 Sheppard on Madness, 5
 Sibson's Medical Anatomy, 3
 Sieveking's Life Assurance, 14
 Smith's (E.) Wasting Diseases of Children, 6
 Clinical Studies, 6
 Smith's (Henry) Surgery of the Rectum, 14
 Smith's (Heywood) Dysmenorrhœa, 6
 Gynæcology, 6
 Smith (Priestley) on Glaucoma, 12
 Southam's Regional Surgery, 10
 Sparks on the Riviera, 10
 Squire's Companion to the Pharmacopœia, 7
 Pharmacopœias of London Hospitals, 7
 Stillé and Maisch's National Dispensatory, 7
 Stocken's Dental Materia Medica and Therapeutics, 12
 Sullivan's Tropical Diseases, 7
 Swain's Surgical Emergencies, 10
 Swayne's Obstetric Aphorisms, 6
 Taft's Operative Dentistry, 12
 Taylor's Medical Jurisprudence, 4
 Poisons in relation to Medical Jurisprudence, 4
 Teale's Dangers to Health, 5
 Thompson's (Sir H.) Calculous Disease, 13
 Diseases of the Prostate, 13
 Diseases of the Urinary Organs, 13
 Lithotomy and Lithotripsy, 13
 Thompson's (Dr. H.) Clinical Lectures, 8
 Thorowgood on Asthma, 8
 on Materia Medica and Therapeutics, 7
 Thudichum's Pathology of the Urine, 14
 Tibbits' Medical and Surgical Electricity, 10
 Map of Motor Points, 10
 Tidy and Woodman's Forensic Medicine, 4
 Tilt's Change of Life, 6
 Uterine Therapeutics, 6
 Tomes' (C. S.) Dental Anatomy, 12
 Tomes' (J. & C. S.) Dental Surgery, 12
 Van Buren on the Genito-Urinary Organs, 14
 Veitch's Handbook for Nurses, 6
 Virchow's Post-mortem Examinations, 4
 Wagstaffe's Human Osteology, 3
 Walker's Ophthalmology, 12
 Waring's Indian Bazaar Medicines, 7
 Practical Therapeutics, 7
 Warner's Guide to Medical Case-Taking, 8
 Waters' (A. T. H.) Diseases of the Chest, 8
 Waters (J. H.) on Fits, 9
 Wells (Spencer) on Ovarian and Uterine Tumours, 6
 West and Duncan's Diseases of Women, 5
 West (S.) How to Examine the Chest, 8
 Whistler's Syphilis of the Larynx, 12
 Whitehead's (J. L.) Climate of the Undercliff, 10
 Whittaker's Primer on the Urine, 14
 Wilks and Moxon's Pathological Anatomy, 4
 Wilson's (Sir E.) Anatomists' Vade-Mecum, 3
 Lectures on Dermatology, 13
 Wilson's (G.) Handbook of Hygiene, 5
 Healthy Life and Dwellings, 5
 Wilson's (W. S.) Ocean as a Health-Resort, 10
 Wise's Davos Platz, 10
 Wolfe's Diseases and Injuries of the Eye, 11
 Yeo's Contagiousness of Pulmonary Consumption, 8

The following CATALOGUES issued by J. & A. CHURCHILL will be forwarded post free on application:—

A. *J. & A. Churchill's General List of about 650 works on Anatomy, Physiology, Hygiene, Midwifery, Materia Medica, Medicine, Surgery, Chemistry, Botany, &c., &c., with a complete Index to their Subjects, for easy reference.*
 N.B.—This List includes B, C, & D.

B. *Selection from J. & A. Churchill's General List, comprising all recent Works published by them on the Art and Science of Medicine.*

C. *J. & A. Churchill's Catalogue of Text Books specially arranged for Students.*

D. *A selected and descriptive List of J. & A. Churchill's Works on Chemistry, Materia Medica, Pharmacy, Botany, Photography, Zoology, the Microscope, and other branches of Science.*

E. *The Half-yearly List of New Works and New Editions published by J. & A. Churchill during the previous six months, together with Particulars of the Periodicals issued from their House.*

[Sent in January and July of each year to every Medical Practitioner in the United Kingdom whose name and address can be ascertained. A large number are also sent to the United States of America, Continental Europe, India, and the Colonies.]

AMERICA.—*J. & A. Churchill being in constant communication with various publishing houses in Boston, New York, and Philadelphia, are able, notwithstanding the absence of international copyright, to conduct negotiations favourable to English Authors.*

LONDON: NEW BURLINGTON STREET.

